arXiv:0712.3678v2 [math.AP] 28 Aug 2008

Numerical approach to L_1-problems with the second order elliptic operators*

Nedžad Limić [†]and Mladen Rogina [‡]

August 28, 2008

Abstract

For a second order differential operator $A(\boldsymbol{x}) = -\nabla a(\boldsymbol{x})\nabla + b'(\boldsymbol{x})\nabla + \nabla\big(\boldsymbol{b}''(\boldsymbol{x})\,\cdot\big)$ on a bounded domain D with the Dirichlet boundary conditions on ∂D there exists the inverse $T(\lambda, A) = (\lambda I + A)^{-1}$ in $L_1(D)$. If μ is a Radon (probability) measure on Borel algebra of subsets of D, then $T(\lambda, A)\mu \in L_p(D), p \in [1, d/(d-1))$. We construct the numerical approximations to $u = T(\lambda, A)\mu$ in two steps. In the first one we construct grid-solutions $\mathbf{u}_n$ and in the second step we embed grid-solutions into the linear space of hat functions $u(n) \in \dot{W}^1_p(D)$. The strong convergence to the original solutions u is established in $L_p(D)$ and the weak convergence in $\dot{W}^1_p(D)$.

AMS subject classification: (2000) 35J20, 35J25, 35J15, 65N06, 65N15

Key words: Elliptic operator, divergence form, difference scheme

1 Introduction

If the tensor-valued function $\{a_{ij}\}_{11}^{dd}$ (diffusion tensor) on a bounded domain $D \subset \mathbb{R}^d$ satisfies the strict ellipticity conditions, then the second order differential operator in divergence form: $A = -\sum_{ij}\partial_i a_{ij}\partial_j + \sum_i b'_i\partial_i + \sum_i \partial_i(b''_i\cdot)$, with homogeneous Dirichlet boundary condition on ∂D, has an inverse A_D^{-1} in terms of an integral operator mapping the Radon measures on Borelian sets $\mathfrak{B}(D)$ into $\dot{W}^1_p(D)$ for each $p \in [1, d/(d-1))$ [BO, LR2]. There are partial results on numerical solutions of the boundary value problem $A_D u = \mu$ [Cl, LR2, LR3]. Here we extend results of [LR3] to a general boundary value problem on $D \subset \mathbb{R}^d$. The operator A_D is discretized by a *system matrix* A_n and numerical solutions are represented by grid-functions $\mathbf{u}_n$. Then $\mathbf{u}_n$ is imbedded into the linear space of hat functions and compared with the solution of the original problem in order to prove convergence. Our particular intention is to construct numerical approximations $\mathbf{u}_n$ with system matrices A_n possessing *compartmental structure*, that is a matrix having positive diagonal elements, non-positive off-diagonal elements and non-negative column sums. Apparently, such matrices are transpose of M-matrices. Because of our determination to look for approximations A_n with the compartmental structure, we have faced a number of non-typical problems in numerical analysis. All of them stem from the construction of the numerical schemes, rather then from the structure of convergence proofs.

Various definitions and auxiliary results which are necessary to prove convergence are given in Sections 2 and 3. In the case of dimension $d = 2$ the proposed numerical

---

*Supported by grant 037-1193086-2771 of the Ministry of Science, Higher Education and Sports, Croatia.

[†] Dept. of Mathematics, University of Zagreb, Bijenička 30, 10002 Zagreb, Croatia, e–mail: nlimic@math.hr

[‡] Dept. of Mathematics, University of Zagreb, Bijenička 30, 10002 Zagreb, Croatia, e–mail: rogina@math.hr

scheme is already on a level of an algorithm and can be straightforwardly applied. The construction of discretizations A_n with the compartmental structure for any dimension is carried out in Section 4. Grid-solutions of discretized problems are imbedded in the linear space of hat functions and the obtained approximate solutions are analyzed from the standpoint of convergence. The convergence in W_2^1-spaces is proved in Section 5 and the convergence in W_p^1-spaces in Section 6. Section 7 provides two examples, demonstrating the efficiency of constructed schemes.

2 Definition of the problem

Elements of $\mathbb{R}^d$ are denoted by $\boldsymbol{x}, \boldsymbol{m}, \boldsymbol{p}$ etc. The Euclidean norm in $\mathbb{R}^d$ is denoted by $|\cdot|$. The only open subsets of $\mathbb{R}^d$ considered in this work, are bounded and connected open sets with Lipshitz boundary [Ma, Ste]. We call a subset of this kind a *domain with Lipshitz boundary.* We denote it by D, and its boundary by ∂D. For a set $S \subset \mathbb{R}^d$ the closure is denoted by $\overline{S}$ and sometimes by $cls(S)$.

The Banach spaces of functions $C^{(k)}(\mathbb{R}^d), C^{(k)}(\overline{D})$ are defined as usually. Their norms are denoted by $\|\cdot\|_\infty^{(k)}$. The closure of functions in $C^{(k)}(\mathbb{R}^d)$ with compact supports determines the subspace $C_0^{(k)}(\mathbb{R}^d)$. The closure of functions in $C(\overline{D})$ with supports in D determines the subspace $\dot{C}(\overline{D})$, and then $\dot{C}^{(k)} = C^{(k)}(\overline{D}) \cap \dot{C}(\overline{D})$. The L_p-spaces as well as Sobolev W_p^1-spaces are defined in a standard way [Ma, Ste]. Their norms are denoted by $\|\cdot\|_p$ and $\|\cdot\|_{p,1}$, respectively. For each p, $1 \le p \le \infty$, the norm of $W_p^1(D)$ is defined by $\|u\|_{p,1} = \big(\|u\|_p^2 + \|\nabla u\|_p^2\big)^{1/2}$, where $\|\, \nabla u \,\|_p = \big(\sum_{j=1}^d \|\, \partial_j u \,\|_p^2\big)^{1/2}$. Because the domain D has Lipshitz boundary, the space $W_\infty^1(D)$ can be realized as the space of continuous functions on $\overline{D}$, for which the first partial derivatives are elements of $L_\infty(D)$. The completion in the norm of $W_p^1(D)$ of functions in $C^{(1)}(D) \cap \dot{C}(\overline{D})$ is denoted by $\dot{W}_p^1(D)$. The linear space $C(\overline{D}) \cap W_p^1(D)$ is dense in $W_p^1(D)$ for all p, $1 \le p \le \infty$, and the closure of $\dot{C}(\overline{D}) \cap W_p^1(D)$ in the norm of $W_p^1(D)$ is equal to $\dot{W}_p^1(D)$ [Ma]. Let X be a Banach space and $X^\dagger$ its dual. Then the value of $f \in X^\dagger$ at $u \in X$ is denoted by $\langle\, f|u \,\rangle$. Let $\mathcal{R}(D)$ be the convex set of positive Radon measures μ on $\mathfrak{B}(D)$. Then $\langle v \,|\, \mu\rangle = \int_D v(\boldsymbol{x})\mu(d\mathbf{x})$ is well defined for $v \in \dot{W}_\infty^1(D)$. We say that a sequence of $\mu_n \in \mathcal{R}(D)$ converges weakly to $\mu \in \mathcal{R}(D)$ if $\lim_n \langle\, v|\mu_n \,\rangle = \langle\, v|\mu \,\rangle$ for each $v \in \dot{C}(\overline{D})$.

Let $\mathbb{1}_S$ be the *indicator* of $S \subset \mathbb{R}^d$, *i.e.* $\mathbb{1}_S(\boldsymbol{x}) = 1$ for $\boldsymbol{x} \in S$, and zero otherwise. We say that f on $\mathbb{R}^d$ is *piecewise continuous with respect to the decomposition* $\mathbb{R}^d = \cup_k D_k$ if there exist a finite collection of L disjoint, Lebesgue measurable subsets $D_k \subset \mathbb{R}^d$, and bounded, uniformly continuous functions on $\mathbb{R}^d$, $\{f_j\}_{j=1}^L$, such that $\mathbb{R}^d = \cup_{k=1}^L D_k$ and $f = \sum_{j=1}^L f_j \mathbb{1}_{D_j}$. If not necessary a part of the terminology such as 'with respect ...' is omitted. A function f_D on $D \subset \mathbb{R}^d$ is piecewise continuous if there exists a piecewise continuous f on $\mathbb{R}^d$ such that $f_D = f|D$. Piecewise constant functions are special cases of piecewise continuous ones.

We consider a 2$^{\text{nd}}$-order elliptic operator on $\mathbb{R}^d$,

$$A(\boldsymbol{x}) = -\sum_{i,j=1}^d \partial_i a_{ij}(\boldsymbol{x})\partial_j \;+ \sum_{j=1}^d b_j'(\boldsymbol{x})\,\partial_j \;+ \sum_{j=1}^d \partial_j\big(b_j''(\boldsymbol{x})\,\cdot\,\big) \;+\; c(\boldsymbol{x}), \tag{1}$$

for which the coefficients must fulfill the following:

Assumption 2.1 *The functions $a_{ij} = a_{ji}$, b_i', b_i'' $(i, j = 1, 2, \ldots, d)$ and c are:*

- *a) piecewise continuous on $\mathbb{R}^d$, $c \ge 0$ and $a_{ij}(\boldsymbol{x})$ converge to constant values as $|\boldsymbol{x}|$ increases,*
- *b) there are positive numbers $\underline{M}, \overline{M}$, $0 < \underline{M} \le \overline{M}$, such that the strict ellipticity*

$$\underline{M}\,|\boldsymbol{x}|^2 \;\le\; \sum_{i,j=1}^d a_{ij}(\boldsymbol{x}) z_i \bar{z}_j \;\le\; \overline{M}\,|\boldsymbol{x}|^2, \quad \boldsymbol{x} \in \mathbb{R}^d \tag{2}$$

holds.

The differential operator $A_0(\boldsymbol{x}) = -\sum_{i,j=1}^{d} \partial_i a_{ij}(\boldsymbol{x})\partial_j$ is called the *main part* of $A(\boldsymbol{x})$.

Let us define a real bilinear form on $W_q^1(D) \times W_p^1(D)$, $1/p + 1/q = 1$, by:

$$\begin{aligned} a(v,u) \;=\; & \sum_{i,j=1}^{d} \int_D a_{ij}(\boldsymbol{x})\, \partial_i v(\boldsymbol{x})\, \partial_j u(\boldsymbol{x})\, d\boldsymbol{x} - \sum_{i=1}^{d} \int_D b_i'(\boldsymbol{x})\, v(\boldsymbol{x})\, \partial_i u(\boldsymbol{x})\, d\boldsymbol{x} \\ & - \sum_{i=1}^{d} \int_D b_i''(\boldsymbol{x})\, \partial_i v(\boldsymbol{x})\, u(\boldsymbol{x})\, d\boldsymbol{x} + \int_D c(\boldsymbol{x})\, v(\boldsymbol{x})\, u(\boldsymbol{x})\, d\boldsymbol{x}. \end{aligned} \tag{3}$$

For a domain D with Lipshitz boundary ∂D and for each pair $v \in \dot{W}_q^1(D)$, $u \in \dot{W}_p^1(D) \cap \{Au \in (\dot{W}_q^1(D))^\dagger\}$, $1 < p < \infty$, the Green formula must be valid,

$$a(v,u) \;=\; \langle\, v \,|\, A\, u \,\rangle.$$

The Green formula is also valid for each pair $v \in \dot{W}_\infty^1(D)$, $u \in \dot{W}_1^1(D) \cap \{Au \in \mathcal{R}(D)\}$.

The boundary value problem, to be studied in this work, is defined by

$$\begin{aligned} \big(\lambda I \,+\, A(\boldsymbol{x})\big)\, u(\boldsymbol{x}) \;&=\; \mu(\boldsymbol{x}), \quad \boldsymbol{x} \in D, \\ u \,\big|\, \partial D \;&=\; 0, \end{aligned} \tag{4}$$

where $\lambda \geq 0$ and D is a domain with Lipshitz boundary. In the case of $D = \mathbb{R}^d$ we suppose that $\lambda > 0$ and the boundary condition in (4) is omitted. The nonhomogeneous term μ is a positive Radon measure for $p \in [1, d/(d-1))$, and $\mu \in W_2^{-1}(D)$ for $p = 2$.

The variational formulation of (4) for a solution $u \in \dot{W}_p^1(D)$, $p \in [1, d/(d-1))$ or $p = 2$, has the form:

$$\lambda(v \,|\, u) \,+\, a(v,u) \;=\; \langle\, v \,|\, \mu \,\rangle, \quad \text{for any } v \in \dot{W}_q^1(D). \tag{5}$$

Solutions of (4) and (5) are called strong and weak solutions, respectively. In the case of a problem on $\mathbb{R}^d$ the variational problem is defined by expression

$$\lambda(v \,|\, u) \,+\, a(v,u) \;=\; \langle\, v \,|\, \mu \,\rangle, \quad \text{for any } v \in W_q^1(\mathbb{R}^d), \tag{6}$$

where for $p = 2$ we have $\mu \in W_2^{-1}(\mathbb{R}^d)$ and for $p \in [1, d/(d-1))$ $\mu \in \mathcal{R}(D)$ with a bounded $D \subset \mathbb{R}^d$.

For the differential operator $H(\boldsymbol{x}) = \lambda I - \sigma^2 \Delta$ on $\mathbb{R}^d$ and $\lambda > 0$ the fundamental solution $(\boldsymbol{x}, \boldsymbol{y}) \mapsto t(\lambda, \boldsymbol{x} - \boldsymbol{y})$ can be represented in terms of the Bessel function $K_\nu, \nu = (d-2)/2$. Let us denote the corresponding operator by $T(\lambda, H)$, i.e. we have $(\lambda I - \sigma^2 \Delta) T(\lambda, H) = I$ on the linear space of continuously differentiable functions with compact supports. Much more, for any $\alpha > 0$ there exists a representation of the operator $T(\lambda, H)^\alpha$ in terms of an integral operator with a positive kernel $t_\alpha(\lambda, \boldsymbol{x} - \boldsymbol{y})$ which is expressed by the Bessel function $K_\nu, \nu = (d - 2\alpha)/2$ [Sh]. The Green function for the differential operator $\lambda I - \sigma^2 \Delta$ on D with the homogeneous Dirichlet boundary conditions at ∂D is denoted by $t(\lambda, H_D, \cdot, \cdot)$ and the corresponding integral operator by $T(\lambda, H_D)$. In this way we have $(\lambda I - \sigma^2 \Delta) T(\lambda, H_D) = I$ on the linear space of continuously differentiable functions on $\overline{D}$. There exist representations of $T(\lambda, H_D)^\alpha$ as integral operators with kernels $t_\alpha(H_D, \cdot, \cdot)$ which are positive on $D \times D$. The equality $(\lambda I + H_D) T(\lambda, D) = I$ enables us to define various closures $H_D = -\lambda I + T(\lambda, H_D)^{-1}$, such as the closure from $\dot{W}_2^1(D)$ onto $W_2^{-1}(D)$, from $\mathfrak{D}(H) = T(\lambda, H_D) L_p(D)$ onto $L_p(D)$ *etc.*

For λ sufficiently large and $\mu \in W_2^{-1}(\mathbb{R}^d)$ solutions to (4)-(6) exist and can be represented by $T(\lambda, H)^{1/2}$ (or $T(\lambda, H_D)^{1/2}$) as described in the following. Let us define the bounded operator on $L_2(\mathbb{R}^d)$ by

$$W \;=\; T(\lambda, H)^{1/2} \Big(\sum_{ij} \partial_i \, (a_{ij} \,-\, \overline{M} \delta_{ij}\,) \partial_j \Big)\, T(\lambda, H)^{1/2}. \tag{7}$$

Then $\|W\|_2 \leq (1-\gamma)$ where $\gamma = \underline{M}/\overline{M} < 1$, so that there exist the operator

$$T(\lambda, A_0) \;=\; T(\lambda, H)^{1/2}\,(I-W)^{-1}\,T(\lambda, H)^{1/2} \tag{8}$$

mapping $W_2^{-1}(\mathbb{R}^d)$ into $W_2^1(\mathbb{R}^d)$ with the norm

$$\|T(\lambda, A_0)\|_{\mathsf{L}(W_2^{-1}, W_2^1)} \leq \frac{\overline{M}}{\underline{M}}\big(\lambda^{-1} + \overline{M}^{-1}\big). \tag{9}$$

From Aronson's inequalities [Ba] we have the following result. The operator $T(\lambda, A_0)$ is an integral operator and its kernel is the fundamental solution of differential operator $\lambda I + A_0(\boldsymbol{x})$ on $\mathbb{R}^d$. If the lower order differential operators in (1) are non-trivial, then $T(\lambda, A)$ exists for λ sufficiently large. Results (8) and (9) are valid for bounded domains as well. We have to replace H with the corresponding H_D and obtain in this way the operators $W_D, T(\lambda, A_D)$. Various closures A, A_D are defined in terms of the constructed operators $T(\lambda, A), T(\lambda, A_0)$, respectively, as in the case of differential operator $-\sigma^2\Delta$.

In the case of $\mu \in \mathcal{R}(D)$ solutions to (4)-(6) also exist. We have the following result [BO, LR2]:

THEOREM **2.1** *Let D be a bounded domain with Lipshitz boundary. For each $p \in [1, d/(d-1))$ there exists a unique weak solution u of (5) belonging to the class $\dot{W}_p^1(D)$ and possessing the following properties:*

(i) There exists a positive number c depending on $\underline{M}, \overline{M}, p, D$, such that the following inequality is valid:

$$\|\,u\,\|_{p,1} \;<\; c\,\mu(D).$$

(ii) If $\{\mu_n : n \in \mathbb{N}\} \subset \mathcal{R}(D)$ converges weakly to a $\mu \in \mathcal{R}(D)$, then the corresponding sequence of weak solutions $\{u_n : n \in \mathbb{N}\} \subset \dot{W}_p^1(D)$, $u_n = A_D^{-1}\mu_n$, converges strongly in $L_p(D)$ to $u = A_D^{-1}\mu$.

This theorem is the theoretical background for construction and analysis of convergence of numerical solutions in $L_1(\mathbb{R}^d)$.

3 Grids and associated functions

Let the orthogonal coordinate system in $\mathbb{R}^d$ be determined by unit vectors $\boldsymbol{e}_i$, and let us define the set G_n by:

$$G_n \;=\; \{\boldsymbol{x} \;=\; h(n)\sum_{l=1}^{d} k_l\,\boldsymbol{e}_l \;:\; k_l \in \mathbb{Z}\}, \tag{10}$$

where $h(n) = 2^{-n}$ is called the grid-step. A grid-step is usually denoted by h and only if necessary by $h(n)$. Elements of G_n are called grid-knots and the constructed sets $G_n, n \in \mathbb{N}$ are called grids. Sometimes we say that G_n discretize $\mathbb{R}^d$. Accordingly, the subgrids $G_n(D) \subset G_n$ defined by $G_n(D) = G_n \cap D$ are called discretizations of D. To each $\boldsymbol{v} \in G_n$ there corresponds a grid-cube $C_n(\mathbf{1}, \boldsymbol{v}) = \prod_1^d [v_j, v_j + h)$, where v_j are coordinates of $\boldsymbol{v} \in G_n$. Cubes $C_n(\mathbf{1}, \boldsymbol{v})$ define a decomposition of $\mathbb{R}^d$ into disjoint sets. Apart from the basic cubes, $C_n(\mathbf{1}, \boldsymbol{v}), \boldsymbol{v} \in G_n$, for constructions we need larger sets. Let $\boldsymbol{p} \in \mathbb{N}^d$. Then

$$C_n(\boldsymbol{p}, \boldsymbol{v}) \;=\; \prod_{i=1}^{d}\,[v_i, v_i + hp_i)$$

are apparently rectangles with "lower left" vertices $\boldsymbol{v}$ and edges of size hp_i. These rectangles define a partition of $\mathbb{R}^d$ as well. The considered cubes $C_n(\mathbf{1}, \boldsymbol{v})$ and rectangles $C_n(\boldsymbol{p}, \boldsymbol{v})$ are semi-closed in the sense that they contain only one of their 2^d vertices.

Basic cubes are defined by their "lower left" vertices. Apart from these basic cubes for our constructions we need closed rectangles,

$$S_n(\boldsymbol{p},\boldsymbol{v}) \;=\; \prod_{i=1}^{d} [v_i - hp_i\,,\; v_i + hp_i], \tag{11}$$

which are defined by central grid-knots $\boldsymbol{v}$. Apparently, $S_n(\boldsymbol{p},\boldsymbol{v})$ is the union of closures of those basic cubes $C_n(\boldsymbol{p},\boldsymbol{x})$ which share the grid-knot $\boldsymbol{v}$.

The grids G_n of (10) are homogeneous with respect to translations in the direction of coordinate axes, i.e $\boldsymbol{x} \in G_n, \boldsymbol{t} = hp_i\boldsymbol{e}_i \Rightarrow \boldsymbol{x}+\boldsymbol{t} \in G_n$ for any $i \in \{1,2,\dots,d\}$ and $p_i \in \mathbb{Z}$. There exist subsets of G_n which are also homogeneous in the defined sense. Let $\boldsymbol{r}_0 \in G_n$ and $\boldsymbol{r} = (r_1, r_2, \dots, r_d) \in \mathbb{N}^d$ be fixed. The set

$$G_n(\boldsymbol{r}_0,\boldsymbol{r}) \;=\; \{\boldsymbol{r}_0 \,+\, h\sum_{l=1}^{d} k_l\, r_l\, \boldsymbol{e}_l \;:\; k_l \in \mathbb{Z}\} \tag{12}$$

is a subset of G_n with the following feature $\boldsymbol{x} \in G_n(\boldsymbol{r}_0,\boldsymbol{r}), \boldsymbol{t} = hp_ir_i\,\boldsymbol{e}_i \Rightarrow \boldsymbol{x}+\boldsymbol{t} \in G_n(\boldsymbol{r}_0,\boldsymbol{r})$. A grid (12) is denoted by $G_n(R)$, where R stands shortly for the $2d$ parameters $\boldsymbol{r}_0, \boldsymbol{r}$.

Let $h_0 = 2^{-n_0}$ for some $n_0 \in \mathbb{N}$, $\boldsymbol{p} \in \mathbb{N}^d$ and let D be a connected set with the structure $D = \cup_{\boldsymbol{v}\in F_n} C_n(\boldsymbol{p},\boldsymbol{v})$, where $F_n \subset G_n$. For the subgrid $G_n(D) = D \cap G_n(R)$ the set $G_n(D)$ is discrete and therefore its interior, closure and boundary are defined indirectly, $int\big(G_n(D)\big) = G_n(D) \cap int(D)$, $cls\big(G_n(D)\big) = G_n(R) \cap \overline{D}$ and $bnd\big(G_n(D)\big)$ is the difference of $cls\big(G_n(D)\big)$ and $int\big(G_n(D)\big)$. Apparently, $int\big(G_n(D)\big) \subseteq G_n(D) \subseteq cls\big(G_n(D)\big)$. Let a finite collection of sets $D_l, l \in \mathfrak{L}$ make a partition of $\mathbb{R}^d$, where each D_l has the structure like the described set D. Then $G(l) = D_l \cap G_n(R)$ make a partition of G_n.

Each $\boldsymbol{x} \in G_n$ can be indexed by $\boldsymbol{m} \in \mathbb{R}^d$, where $\boldsymbol{x} = h\boldsymbol{m}$. Similarly, we index grid-knots of $G_n(\boldsymbol{r}_0,\boldsymbol{r})$ by those $\boldsymbol{m} \in \mathbb{Z}^d$ for which there holds $\boldsymbol{x} = \boldsymbol{r}_0 + h\sum_l m_l r_l \boldsymbol{e}_l$. Therefore, we define the sets $I_n = \mathbb{Z}^d$ and $I_n(R) \subset I_n$, indexing the grid-knots of G_n and $G_n(R)$. In this work frequently utilized pairs of grids and their index sets are $G_n,\ I_n;\ G_n(R),\ I_n(R);\ G_n(l),\ I_n(l);\ G_n(R,D), I_n(R,D)$.

The *shift operator* $Z(\boldsymbol{x}), \boldsymbol{x} \in \mathbb{R}^d$, acting on functions $f : \mathbb{R}^d \mapsto \mathbb{R}$, is defined by $\big(Z(\boldsymbol{x})f\big)(\boldsymbol{x}) = f(\boldsymbol{x}+\boldsymbol{z})$. Similarly we define the discretized shift operator by $\big(Z_n(r,i)\mathbf{u}_n\big)_{\boldsymbol{k}} = (\mathbf{u}_n)_{\boldsymbol{l}}$, where $\boldsymbol{l} = \boldsymbol{k} + rh\boldsymbol{e}_i$.

Discretization of differential operators. With respect to a grid step h, the partial derivatives of $u \in C^{(1)}(\mathbb{R}^d)$ are discretized by forward/backward finite difference operators in the usual way,

$$\begin{array}{rcl} \eth_i(t)u(\boldsymbol{x}) & = & \frac{1}{t}\big(u(\boldsymbol{x}+t\boldsymbol{e}_i) \,-\, u(\boldsymbol{x})\big), \\ \widehat{\eth}_i(t)u(\boldsymbol{x}) & = & \frac{1}{t}\big(u(\boldsymbol{x}) \,-\, u(\boldsymbol{x}-t\boldsymbol{e}_i)\big), \end{array} \qquad \boldsymbol{x} \in \mathbb{R}^d,\ t \neq 0. \tag{13}$$

Let $r \in \mathbb{Z}\backslash\{0\}$. Discretizations of the functions $\partial_i u$ on G_n, denoted by $U_i(r)\mathbf{u}_n, V_i(r)\mathbf{u}_n$, are defined by:

$$\big(U_i(r)\,\mathbf{u}_n\big)_{\boldsymbol{m}} \;=\; \eth_i\big(rh\big)\,u(\boldsymbol{x}_{\boldsymbol{m}}), \quad \big(V_i(r)\,\mathbf{u}_n\big)_{\boldsymbol{m}} \;=\; \widehat{\eth}_i\big(rh\big)\,u(\boldsymbol{x}_{\boldsymbol{m}}),$$

where $\boldsymbol{x} \in G_n$. Then

$$\begin{array}{rcl} U_i(r) & = & (rh)^{-1}(Z_n(r,i) \,-\, I), \\ V_i(r) & = & (rh)^{-1}\big(I - Z_n(-r,i)\big) \;=\; U_i(-r) \;=\; -\,U_i(r)^T. \end{array} \tag{14}$$

Therefore we have $U_i(-r) = U_i(r)\,Z_n(-r,i) = Z_n(-r,i)\,U_i(r)$, and similarly for $V_i(r)$. In the case of $r = 1$ we use a short notations U_i, V_i.

In accordance with the previous terminology, we say that $\partial_i, \sum_{ij}\partial_i a_{ij}\partial_j$ *etc.* are differential operators on $\mathbb{R}^d$ or D. We say that their discretizations are defined on G_n or $G_n(D)$. In particular, discretizations of the differential operator (1) are denoted by A_n. Naturally, matrices A_n are the main object in this work.

3.1 Relations between $l(G_n(R))$ and W_2^1-spaces

The discretization of a function $u \in C(\mathbb{R}^d)$ on G_n is denoted by $\mathbf{u}_n$ and defined by values at grid-knots, $(\mathbf{u}_n)_{\boldsymbol{m}} = u(\boldsymbol{x}_{\boldsymbol{m}})$ where $\boldsymbol{x}_{\boldsymbol{m}} = (m_1 h, m_2 h, \ldots, m_d h) \in G_n$, and $\boldsymbol{m} = (m_1, m_2, \ldots, m_d)$ is a multi-index. The function $\mathbf{u}_n$ is usually called a grid-function. We denote the linear spaces of grid-functions by $l(G_n)$ or $l(G_n(D))$. The linear space of grid-functions on $G_n(R)$ with finite supports is denoted by $l_0(G_n(R))$. Elements of $l(G_n)$ are also called columns. The corresponding L_p-spaces are denoted by $l_p(G_n)$ or $l_p(G_n(D))$, and their norms by $\|\cdot\|_p$. The duality pairing of $\mathbf{v} \in l_q(G_n)$ and $\mathbf{u} \in l_p(G_n)$ is denoted by $\langle \mathbf{v}|\mathbf{u} \rangle$. The scalar product in $l_2(G_n)$ is denoted by $\langle \cdot|\cdot \rangle$ and sometimes by $(\cdot|\cdot)$. The norm of $l_p(G_n(R))$ is denoted by $\|\cdot\|_{Rp}$. For $p \in [1, \infty)$ this norm is defined by:

$$\|\mathbf{u}\|_{Rp} = \Big[\mathrm{vol}(R) \sum_{\boldsymbol{k} \in I_n(R)} |u_{\boldsymbol{k}}|^p \Big]^{1/p},$$

where $\mathrm{vol}(R) = \prod_{i=1}^d r_i$. Finally, for $p = \infty$ we have $\|\mathbf{u}\|_{R\infty} = \sup\{|u_{\boldsymbol{k}}| : \boldsymbol{k} \in I_n(R)\}$.

Let us define the quadratic functional on $l(G_n)$ by $q(\mathbf{u}) = \sum_i^d \|U_i \mathbf{u}\|_2^2$ and $q_R(\mathbf{u}) = \mathrm{vol}(R) \sum_i^d \|U_i(r_i)\mathbf{u}\|_{R2}^2$ on $l(G_n(R))$. It is understood $q_R = q$ for $G_n(R) = G_n$. A discrete analog of W_2^1-spaces is the linear spaces of those $\mathbf{u}_n \in l(G_n(R))$ for which the norm $\|\cdot\|_{R2,1}$:

$$\|\mathbf{u}\|_{R2,1}^2 = \|\mathbf{u}\|_{R2}^2 + q_R(\mathbf{u}), \tag{15}$$

is finite. This space is denoted by $w_2^1(G_n(R))$. By convention $\|\cdot\|_{2,1} = \|\cdot\|_{R2,1}$ for $r_i = 1$. The subspace of grid-functions $\mathbf{u} \in w_2^1(G_n(R))$ for which $\mathbf{u}_n = \mathbb{1}_{G_n(D)}\mathbf{u}_n$ is denoted by $w_2^1(G_n(R, D))$. Hence, $w_2^1(G_n(R, D))$ for $\boldsymbol{r} = \mathbf{1}$ is denoted by $w_2^1(G_n(D))$. For problems on bounded domains we need a discrete version of the Poincaré inequality which is formulated as follows:

LEMMA **3.1** *Let D be bounded. Then the norms $\|\cdot\|_{2,1}$ and $q_R(\cdot)^{1/2}$ are equivalent in $w_2^1(G_n(R, D))$,*

$$q_R(\cdot)^{1/2} \geq \beta \|\cdot\|_{R2,1},$$

where β is independent of n.

An element (column) $\mathbf{u}_n \in l(G_n)$ can be associated to a continuous function on $\mathbb{R}^d$ in various ways. Here is utilized a mapping $l(G_n) \mapsto C(\mathbb{R}^d)$ which is defined in terms of hat functions. Let χ be the canonical hat function on $\mathbb{R}$, centered at the origin and having the support $[-1, 1]$. Then $z \mapsto \phi(h, x, z) = \chi(h^{-1}(z - hx))$ is the hat function on $\mathbb{R}$, centered at $x \in \mathbb{R}$ with support $[x - h, x + h]$. The functions $\boldsymbol{z} \mapsto \phi_{\boldsymbol{k}}(\boldsymbol{z}) = \prod_{i=1}^d \phi(h, x_i, z_i), x_i = hk_i$, define d-dimensional hat functions with supports $S_n(\mathbf{1}, \boldsymbol{x}) = \prod_i [x_i - h, x_i + h]$. The functions $\phi_{\boldsymbol{k}}(\cdot) \in G_n$, span a linear space, denoted by $E_n(\mathbb{R}^d)$. Let $\mathbf{u}_n \in l(G_n)$ have the entries $u_{n\boldsymbol{k}} = (\mathbf{u}_n)_{\boldsymbol{k}}$. Then the function $u(n) = \sum_{\boldsymbol{k} \in I_n} u_{n\boldsymbol{k}} \phi_{\boldsymbol{k}}$ belongs to $E_n(\mathbb{R}^d)$ and defines imbedding of grid-functions into the space of continuous functions. We denote the corresponding mapping by $\Phi_n : l(G_n) \mapsto E_n(\mathbb{R}^d)$. Obviously that there exists $\Phi_n^{-1} : E_n(\mathbb{R}^d) \mapsto l(G_n)$ and the spaces $l(G_n)$ and $E_n(\mathbb{R}^d)$ are isomorphic with respect to the pair of mappings Φ_n, Φ_n^{-1}. It is clear that $E_n(\mathbb{R}^d) \subset E_{n+1}(\mathbb{R}^d)$ and the space of functions $\cup_n E_n(\mathbb{R}^d)$ is dense in $L_p(\mathbb{R}^d), p \in [1, \infty)$, as well as in $\dot{C}(\mathbb{R}^d)$. Let us mention that $\sum_{\boldsymbol{k}} \phi_{\boldsymbol{k}} = 1$ on $\mathbb{R}^d$.

Now we consider another collection of basis functions. To each $\boldsymbol{x} = h\boldsymbol{k} \in G_n(R)$ there is associated a d-dimensional hat function

$$\psi_{\boldsymbol{k}}(\boldsymbol{x}) = \prod_{i=1}^d \chi\left(\frac{x_i - hk_i}{hr_i} \right),$$

obviously, with the support $S_n(\boldsymbol{r}, \boldsymbol{x}) = \prod_i [x_i - r_i h, x_i + r_i h]$. They span a linear space denoted by $E_n(R, \mathbb{R}^d)$. Again we have $\sum_{\boldsymbol{k}} \psi_{\boldsymbol{k}} = 1$ on $\mathbb{R}^d$. The mappings Φ_n, Φ_n^{-1}

cannot be applied to elements of $l(G_n(R))$ and $E_n(R,\mathbb{R}^d)$, respectively. Therefore we define restrictions $\Phi_n(R) : l(G_n(R)) \to E_n(R,\mathbb{R}^d)$ and the corresponding inverse mapping $\Phi_n^{-1}(R)$ by the following expression:

$$u(n) \;=\; \Phi_n(R)\,\mathbf{u}_n \;=\; \sum_{\boldsymbol{k}} \big(\mathbf{u}_n\big)_{\boldsymbol{k}}\,\psi_{\boldsymbol{k}}. \tag{16}$$

If we have to underline that $u(n)$ is related to a particular set of parameters R then we use an extended denotation $u(R,n)$. For two functions $v(n), u(n)$ we have $(v(n)|u(n)) = h^d \mathrm{vol}(R) \sum_{\boldsymbol{kl}} s_{\boldsymbol{kl}} v_{\boldsymbol{k}} u_{\boldsymbol{k}}$ where $s_{\boldsymbol{kl}} = \|\psi_{\boldsymbol{k}}\|_1^{-1}(\psi_{\boldsymbol{k}}|\psi_{\boldsymbol{l}})$. Let us notice that $\sum_{\boldsymbol{l}} s_{\boldsymbol{kl}} = 1$.

We cannot compare directly columns $\mathbf{u}_n$ with various n. An indirect comparison can be made by using $u(n) = \Phi_n(R)\mathbf{u}_n \in \cup_n E_n(R,\mathbb{R}^d)$. To compare $U_i(r_i)\mathbf{u}_n$ and $\partial_i u(n)$ we need an additional expression. Let $\mathbf{u}$ and $u(n)$ be related by (16) and $\dot{s}_{\boldsymbol{kl}}(i) = h^{-d}(\partial_i\psi_{\boldsymbol{k}}|\partial_i\psi_{\boldsymbol{l}})$. Then, for a homogeneous grid $G_n(R), \boldsymbol{r} \in \mathbb{N}^d$, there must hold

$$\sum_{\boldsymbol{kl}} v_{\boldsymbol{k}} u_{\boldsymbol{l}}\, \dot{s}_{\boldsymbol{kl}}(i) \;=\; -\frac{1}{2} \sum_{\boldsymbol{k}\,\boldsymbol{r}'\,r_i} \big(v_{\boldsymbol{k}+r_i\boldsymbol{e}_i} \,-\, v_{\boldsymbol{k}}\big)\big(u_{\boldsymbol{k}+\boldsymbol{r}'+r_i\boldsymbol{e}_i} \,-\, u_{\boldsymbol{k}+\boldsymbol{r}'}\big)\, s'_{\boldsymbol{0r}'}\, \dot{s}_{0r_i}, \tag{17}$$

where $\boldsymbol{r}' = (r_1, r_2, \ldots, r_{i-1}, 0, r_{i+1}, \ldots, r_d)$, $\dot{s}_{0r} = (\partial_i\psi_0|\partial_i\psi_r)$ and $s'_{\boldsymbol{0r}'}$ is the $(d-1)$-dimensional quantity $s_{\boldsymbol{kl}}$. Then (17) follows from $\sum_{\boldsymbol{l}} s_{\boldsymbol{kl}} = 1$ and consequently $\sum_{\boldsymbol{l}} \dot{s}_{\boldsymbol{kl}} = 0$, after the sum is carried out over any partial component l_i of the index $\boldsymbol{l}$. Thus we have

$$(\partial_i v(n)\,|\,\partial_i u(n)) \;=\; \|\psi_{\boldsymbol{k}}\|_1 \sum_{m_i\boldsymbol{k}'\boldsymbol{l}'} s_{\boldsymbol{k}'\boldsymbol{l}'}\, \big(U_i(r_i)\mathbf{v}\big)_{m_i\boldsymbol{k}'}\, \big(U_i(r_i)\mathbf{u}\big)_{m_i\boldsymbol{l}'}, \tag{18}$$

where the indices are defined by $\boldsymbol{k}' = (k_1, \ldots, k_{i-1}, k_{i+1}, \ldots, k_d)$ and analogously $\boldsymbol{l}'$.

LEMMA **3.2** *Let sequences of functions $v(n), u(n), n \in \mathbb{N}$, be defined by (16). Then*

(i) $$\Big|\textstyle\sum_{i=1}^d \big(\partial_i v(n)\,|\,\partial_i u(n)\big)\Big| \;\le\; h^d\, q_R(\mathbf{v})^{1/2}\, q_R(\mathbf{u})^{1/2}.$$

(ii) $$\Big|\big(\partial_i v(n)\,|\,\partial_j u(n)\big) \,-\, h^d\,\mathrm{vol}(R) \textstyle\sum_{\boldsymbol{k}\in I_n(R)} \big(U_i(r_i)\mathbf{v}\big)_{\boldsymbol{k}}\, \big(U_j(r_j)\mathbf{u}\big)_{\boldsymbol{k}}\Big|$$
$$\le\; h^d\, \min \left\{ \begin{array}{ll} \boldsymbol{|}U_i(r_i)\mathbf{v}\boldsymbol{|}_{R2} & \sup\big\{\boldsymbol{|}(Z(w,j)-I)\,U_j(r_j)\mathbf{u}\boldsymbol{|}_{R2} \,:\, |w| \le r_j h\big\} \\ \boldsymbol{|}U_j(r_j)\mathbf{u}\boldsymbol{|}_{R2} & \sup\big\{\boldsymbol{|}(Z(w,i)-I)\,U_i(r_i)\mathbf{v}\boldsymbol{|}_{R2} \,:\, |w| \le r_i h\big\}. \end{array}\right.$$

PROOF: After applying the CSB-inequality to (18) and using $\sum_{\boldsymbol{l}} s_{\boldsymbol{kl}} = 1$ we get (i). Assertion (ii) is proved here for $i = j = 1$. In this proof $\partial = \partial_1$. By (18) we can straightforwardly calculate

$$\begin{aligned}(\partial v(n)\,|\,\partial u(n)) \;&=\; \|\psi_{\boldsymbol{k}}\|_1 \sum_{\boldsymbol{k},\boldsymbol{r}'} s_{\boldsymbol{0r}'}\, \big(U_i(r_i)\mathbf{v}\big)_{\boldsymbol{k}}\, \big(U_i(r_i)\mathbf{u}\big)_{\boldsymbol{k}+\boldsymbol{r}'} \\ &=\; h^d\, \big(U_i(r_i)\mathbf{v}\big|U_i(r_i)\mathbf{u}\big)_R \,+\, \delta(n),\end{aligned}$$

where

$$\delta(n) \;=\; \|\psi_{\boldsymbol{k}}\|_1 \sum_{\boldsymbol{k},\boldsymbol{r}'} s_{\boldsymbol{0r}'}\, \big(U_i(r_i)\mathbf{v}\big)_{\boldsymbol{k}} \Big[\big(U_i(r_i)\mathbf{u}\big)_{\boldsymbol{k}+\boldsymbol{r}'} \,-\, \big(U_i(r_i)\mathbf{u}\big)_{\boldsymbol{k}}\Big].$$

By the CBS inequality the error term $\delta(n)$ can be estimated as in Assertion (ii).**QED**

LEMMA **3.3** *Let $G_n(R)$ be a homogeneous subgrid defined by (12). There exists $\sigma^2 \in (0,1)$ such that*

$$(1 \,-\, \sigma^2)\, h^d\, \boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{R2}^2 \;\le\; \|\, u(n)\,\|_2^2 \;\le\; h^d\, \boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{R2}^2$$

uniformly with respect to $n \in \mathbb{N}$.

Proof: Let us consider first the one-dimensional case. The grid $G_n(R)$ consists of points $x_k = hrk \in \mathbb{R}, k \in \mathbb{Z}$, and $E_n(R,\mathbb{R})$ is spanned by the hat functions ψ_k centered at x_k with the supports $[-hr + x_k, x_k + hr]$. We define the matrix $S(1)$ with entries:

$$s_{kl} \;=\; \frac{1}{hr}\,(\psi_k|\psi_l) \;=\; \begin{cases} (2/3) & \text{for} \quad k \;=\; l, \\ (1/6) & \text{for} \quad k \;=\; l \pm r. \end{cases}$$

Obviously we have $S(1) = I - (1/3)A$, where the matrix A has the structure $A = I + (1/2)(I_+ + I_-)$ and $I_\pm$ are the first upper and lower off-diagonals. It is well known [Str] that A has a purely continuous spectrum in $[0,2]$ so that $S(1)$ has the spectrum equal $[1/3, 1]$. Therefore

$$\|u(n)\|_2^2 \;=\; hvol(R)\sum_{kl} s_{kl}\,u_k u_l \geq \frac{1}{3}\,hvol(R)\sum_k u_k^2 \;=\; \frac{1}{3}\,h\,\bracevert\mathbf{u}\bracevert_{R2}^2.$$

Hence, we have here $1 - \sigma^2 = 1/3$.

In order to generalize this proof to d-dimensional case we proceed as follows. The symmetric matrix $S(d)$ with entries $s_{\boldsymbol{kl}}$ can be represented as the outer product of d matrices $S(1)$ with entries as in the first part of proof. Therefore its spectrum is $Sp(S(d)) = \prod_{i=1}^d Sp(S(1))$. According to the first part of proof the matrix $S(1)$ has its spectrum in the interval $[1/3, 1]$, implying $Sp(d) \geq 3^{-d}$. Hence, with $\sigma^2 = 1 - 3^{-d}$ we have

$$\mathrm{vol}(R)\sum_{\boldsymbol{kl}\in I_n(R)} s_{\boldsymbol{kl}}\,u_{\boldsymbol{k}}\,u_{\boldsymbol{l}} \;\geq\; (1-\sigma^2)\,\bracevert\mathbf{u}\bracevert_{R2}^2, \tag{19}$$

providing us with a proof of the general case. **QED**

THEOREM **3.1** *Let $u(n) = \Phi_n(R)\mathbf{u}_n$. There exists $\sigma^2 \in (0,1)$, independent of n, such that*

$$(1-\sigma^2)\,h^d\,\bracevert\mathbf{u}_n\bracevert_{R2,1}^2 \;\leq\; \|u(R,n)\|_{2,1}^2 \;\leq\; h^d\,\bracevert\mathbf{u}_n\bracevert_{R2,1}^2.$$

Proof: It is sufficient to prove the first double inequality. The estimates from above are obvious. To get the estimates from bellow it suffices to consider $\partial_i u$. From Expression (18) we have

$$\|\partial_i u(n)\|_2^2 \;=\; \|\psi_{\boldsymbol{k}}\|_1 \sum_m \sum_{\boldsymbol{k'l'}} s_{\boldsymbol{k'l'}}\,\big(U_i(r_i)\mathbf{u}\big)_{m\boldsymbol{k'}}\,\big(U_i(r_i)\mathbf{u}\big)_{m\boldsymbol{l'}}.$$

Then after applying (19) to the inner sum we get

$$\|\partial u(n)\|_2^2 \;\geq\; (1-\sigma^2)\,\|\psi_{\boldsymbol{k}}\|_1 \sum_{m,\boldsymbol{k'}} \big(U_i(r_i)\mathbf{u}\big)_{m\boldsymbol{k'}}^2 \;=\; (1-\sigma^2)\,h^d\,\bracevert U_i(r_i)\mathbf{u}\bracevert_{R2}^2,$$

from where follows the estimate from bellow. **QED**

An element $u \in W_2^1(\mathbb{R}^d)$ does not necessary belong to $E_n(R,\mathbb{R}^d)$. In order to approximate u with elements of $E_n(R,\mathbb{R}^d)$ we define:

$$\hat{u}(n) \;=\; \sum_{\boldsymbol{k}\in I_n(R)} \|\psi_{\boldsymbol{k}}\|_1^{-1}\,(\psi_{\boldsymbol{k}}|u)\,\psi_{\boldsymbol{k}}. \tag{20}$$

The numbers $\|\psi_{\boldsymbol{k}}\|_1^{-1}(\psi_{\boldsymbol{k}}|u)$ are called the *Fourier coefficients* of u.

The basic result for our proof of convergence of approximate solutions is formulated in terms of the quantity $\Gamma_p(\boldsymbol{w}, u)$ defined by:

$$\Gamma_p(\boldsymbol{w}, u) \;=\; \|\,(Z(\boldsymbol{w}) \;-\; I)\,u\,\|_p$$

The kernels

$$\omega_n(\boldsymbol{x},\boldsymbol{y}) \;=\; \sum_{\boldsymbol{k}} \frac{1}{\|\psi_{\boldsymbol{k}}\|_1}\,\psi_{\boldsymbol{k}}(\boldsymbol{x})\,\psi_{\boldsymbol{k}}(\boldsymbol{y}) \tag{21}$$

define an integral operator which is denoted by K_n. Actually, the kernels ω_n define a δ-sequence of functions on $\mathbb{R}^d \times \mathbb{R}^d$ and K_n converge strongly in L_p-spaces to unity:

COROLLARY **3.1** *Let $p \in [1,\infty]$. Then*

- *(i)* $\|K_n\|_p \leq 1$.
- *(ii) There is a positive number $\kappa(R)$, independent of n, such that $\|(I-K_n)u\|_p \leq \kappa(R)\sup\{\Gamma_p(\boldsymbol{w},u) \,:\, |w_i| \leq hr_i\}$.*
- *(iii) The operator $K_n \in \mathsf{L}(L_2(\mathbb{R}^d), L_2(\mathbb{R}^d))$ has the spectrum equal $Sp(K_n) = \{0\} \cup [3^{-d}, 1]$.*

Proof: Only (iii) has to be proved. The symmetric operator K_n is reduced by $E_n(R,\mathbb{R}^d)$ and represented by an integral operator with the kernel (21). It is zero operator in the orthogonal complement $E_n(R,\mathbb{R}^d)^{\perp}$. With respect to the mapping $\Phi_n(R)$ the operator K_n is mapped to the symmetric matrix $\hat{K_n} = \Phi_n(R)^{-1}K_n\Phi_n(R) = S(d)$ in $l_2(I_n(R))$. **QED**

THEOREM **3.2** *Let $v,u \in W_2^1(\mathbb{R}^d)$ and $\hat{u}(n), \hat{v}(n)$ be defined by (20). Then*

$$\big|(\hat{v}(n)|\hat{u}(n)) \;-\; (v|u)\big| \;\leq\; c(R)\,\min\left\{\begin{array}{l} \|\,u\,\|_2\ \sup_{|\boldsymbol{w}|\leq h|\boldsymbol{r}|}\ \Gamma_2(\boldsymbol{w},v), \\ \|\,v\,\|_2\ \sup_{|\boldsymbol{w}|\leq h\boldsymbol{r}|}\ \Gamma_2(\boldsymbol{w},u), \end{array}\right.$$

$$\begin{aligned} &\big|(\partial_i\hat{v}(n)\,|\,\partial_j\hat{u}(n)) \;-\; (\,\partial_i v\,|\,\partial_j u\,)\big| \\ &\leq\; c(R)\,\min\left\{\begin{array}{l} \|\partial_i v\|_2\Big[\|\partial_j u - \exists_j u\|_2 \;+\; \sup\limits_{|\boldsymbol{w}|\leq h\boldsymbol{r}|}\ \Gamma_2(\boldsymbol{w},\partial_j u)\Big], \\ \|\partial_j u\|_2\Big[\|\partial_i v - \exists_i v\|_2 \;+\; \sup\limits_{|\boldsymbol{w}|\leq h\boldsymbol{r}|}\ \Gamma_2(\boldsymbol{w},\partial_i v)\Big], \end{array}\right. \end{aligned}$$

where $c(R)$ is n-independent.

A proof of this theorem is rather technical. For instance, in order to prove the second inequality one has to use a sequence of replacements: $(\partial_i\hat{v}(n)|\partial_j\hat{u}(n)) \to (\exists_i(h)v|K_n\exists_j(h)u) \to (\exists_i(h)v|K_n\partial_j u) \to (\partial_i v|K_n\partial_j u) \to (\partial_i v|\partial_j u)$. Each replacement gives rise to an error. The sum of errors can be estimated by an expression as given in the second inequality of assertion.

3.2 Imbedding of $l(G_n(R))$ into W_2^{-1}-spaces

Beside the functions $\psi_{\boldsymbol{k}}$ we consider the functions defined by:

$$\chi_{\boldsymbol{k}i+}(\boldsymbol{x}) \;=\; \mathbb{1}_{[k_i,k_i+r_ih)}(x_i)\prod_{j\neq i}\psi_{k_j}(x_j), \quad \chi_{\boldsymbol{k}i-}(\boldsymbol{x}) \;=\; \chi_{\boldsymbol{k}-r_i\boldsymbol{e}_i i+}(\boldsymbol{x})$$

for all the possible $i = 1,2,\ldots,d$, and the linear space $F_n(R,\mathbb{R}^d)$ spanned by the defined functions $\chi_{\boldsymbol{k}i\pm}$. The integral operators $K_n^{(i)}(\chi,\chi)$ with the respective kernels

$$\omega_n^{(i)}(\boldsymbol{x},\boldsymbol{y}) \;=\; \sum_{\boldsymbol{k}}\frac{1}{\|\chi_{\boldsymbol{k}i+}\|_1}\,\chi_{\boldsymbol{k}i+}(\boldsymbol{x})\,\chi_{\boldsymbol{k}i+}(\boldsymbol{y}) \;=\; \sum_{\boldsymbol{k}}\frac{1}{\|\chi_{\boldsymbol{k}i+}\|_1}\,\chi_{\boldsymbol{k}i-}(\boldsymbol{x})\,\chi_{\boldsymbol{k}i-}(\boldsymbol{y})$$

have properties similar to the integral operators K_n of Corollary 3.1. The same is valid for the non-symmetric integral operators $K_n^{(i)}(\psi,\chi)$, and their adjoints $K_n^{(i)}(\chi,\psi)$, where the kernel of $K_n^{(i)}(\psi,\chi)$ is defined by $(\boldsymbol{x},\boldsymbol{y}) \mapsto (h^d\mathrm{vol}(R))^{-1}\sum_{\boldsymbol{k}}\psi_{\boldsymbol{k}}(\boldsymbol{x})\boldsymbol{\chi}_{\boldsymbol{k}i+}(\boldsymbol{y})$.

LEMMA **3.4** *The operators $K_n^{(i)}(\chi,\chi), K_n^{(i)}(\psi,\chi), K_n^{(i)}(\psi,\chi)^{\dagger}$ have properties (i) and (ii) of Corollary 3.1. The spectra of operators $K_n^{(i)}(\chi,\chi) \in \mathsf{L}(L_2(\mathbb{R}^d), L_2(\mathbb{R}^d))$ consist of 0 and an interval $[\kappa,1]$ with certain $\kappa \in (0,1)$.*

An element $\mu \in W_2^{-1}(\mathbb{R}^d)$ is represented as $\mu = f_0 + \sum_{i=1}^d \partial_i f_i$ with $f_i \in L_2(\mathbb{R}^d)$ with the norm $\|\mu\|_{2,-1} = (\sum_i \|f_i\|_2^2)^{1/2}$. Its discretizations are defined by grid-functions $\boldsymbol{\mu}_n$ with the components:

$$\mu_{\boldsymbol{k}} \;=\; \frac{1}{\|\psi_{\boldsymbol{k}}\|_1}\,(\psi_{\boldsymbol{k}}|f_0) \;-\; \frac{1}{\|\psi_{\boldsymbol{k}}\|_1}\,\sum_{i=1}^{d}(\partial_i\,\psi_{\boldsymbol{k}}|f_i). \tag{22}$$

Therefore we can write:

$$\langle\, u(n) \,|\, \mu \,\rangle \;=\; h^d \,\langle\, \mathbf{u}_n \,|\, \boldsymbol{\mu}_n \,\rangle_R \;=\; \|\psi_{\boldsymbol{k}}\|_1 \sum_{\boldsymbol{k}\in I_n(R)} u_{\boldsymbol{k}}\mu_{\boldsymbol{k}}. \tag{23}$$

The expression

$$\partial_i\psi_{\boldsymbol{k}} \;=\; \frac{1}{r_i h}\Big[\chi_{\boldsymbol{k}i-} - \chi_{\boldsymbol{k}i+}\Big], \quad a.e. \text{ on } \mathbb{R}^d,$$

enables us to rewrite the components of $\boldsymbol{\mu}_n$ in (22) in terms of the Fourier coefficients $\hat{f}_{i\boldsymbol{k}\pm} = \|\psi_{\boldsymbol{k}}\|_1^{-1}(\chi_{\boldsymbol{k}i\pm}|f_i)$ of functions f_i. The corresponding grid-functions $\hat{\mathbf{f}}_{in\pm}$ are imbedded into the space $F_n(R,\mathbb{R}^d)$ by the mappings $\check{\mathbf{f}}_{in\pm} \mapsto \check{f}_{i\pm}(n) = K_n^{(i)}(\chi,\chi)f_i$ which are completely analogous to the mapping $\hat{\mathbf{f}}_0 \mapsto \hat{f}_0(n) = K_n f_0$ of Corollary 3.1. Now we can get the following useful expression:

$$\langle\, u(n) \,|\, \mu \,\rangle \;=\; h^d \Big[(\mathbf{u}_n|\hat{\mathbf{f}}_{0n})_R \;-\; \sum_{i=1}^{d} (U_i(r_i)\mathbf{u}_n|\check{\mathbf{f}}_{in+})\Big]. \tag{24}$$

Therefore, by (14) we can represent elements $\boldsymbol{\mu} \in l(G_n(R))$ in the following way:

$$\boldsymbol{\mu}_n \;=\; \hat{\mathbf{f}}_{0n} \;-\; \sum_{i=1}^{d} V_i(r_i)\check{\mathbf{f}}_{in+}. \tag{25}$$

From Expressions (23) and (24) we have $|\langle\, \mathbf{u}_n|\boldsymbol{\mu}_n\,\rangle| \le \boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{2,1}(\boldsymbol{|}\hat{\mathbf{f}}_0\boldsymbol{|}_2^2 + \boldsymbol{|}\check{\mathbf{f}}_{in+}\boldsymbol{|}_2^2)^{1/2}$, implying a natural definition:

$$\boldsymbol{|}\boldsymbol{\mu}_n\boldsymbol{|}_{2,-1}^2 \;=\; \sum_{i=1}^{d} \boldsymbol{|}\hat{\mathbf{f}}_0\boldsymbol{|}_2^2 + \boldsymbol{|}\check{\mathbf{f}}_{in+}\boldsymbol{|}_2^2.$$

For the sake of a concise writing of final results we denote here K_n by $K_n^{(0)}$.

LEMMA **3.5** *For each $\mu \in W_2^{-1}(\mathbb{R}^d)$ and the corresponding discretizations $\boldsymbol{\mu}_n$ defined by (25) the following is valid:*

$$h^d \,\boldsymbol{|}\boldsymbol{\mu}_n\boldsymbol{|}_{2,-1}^2 \;=\; \sum_{i=0}^{d} (f_i \,|\, K_n^{(i)}(\chi,\chi)f_i) \;\le\; \|\mu\|_{2,-1}^2,$$

and

$$0 \;\le\; \|\mu\|_{2,-1}^2 \;-\; h^d \,\boldsymbol{|}\boldsymbol{\mu}_n\boldsymbol{|}_{2,-1}^2 \;=\; \sum_{i=0}^{d} (f_i \,|\, (I - K_n^{(i)}(\chi,\chi))f_i) \;\to\; 0,$$

as $n \to \infty$.

3.3 Imbedding into W_p^1-spaces

Theorem 3.1 can be partially generalized.

LEMMA **3.6** *The following assertions are valid:*

(i) If $p \in [1,\infty]$ and $\mathbf{u}_n \in l_p(G_n(R))$ then:

$$\|\, u(n) \,\|_{p,1} \;\le\; h^{d/p}\,\boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{Rp,1}.$$

(ii) If $p \in [1,\infty]$ and $u \in W_p^1(\mathbb{R}^d)$ then:

$$\|\hat{u}(n)\|_{p,1} \;\le\; h^{d/p}\,\boldsymbol{|}\hat{\mathbf{u}}_n\boldsymbol{|}_{Rp,1} \;\le\; \|u\|_{p,1}.$$

(iii) Let $u = u^+ + u^-$*, where* $u^+ = \max\{u, 0\}, u^- = \min\{u, 0\}$*. Then*

$$h^d \,\boldsymbol{|}\hat{\mathbf{u}}_n\boldsymbol{|}_{R1} \;\le\; \|\, u \,\|_1 \;=\; \|\hat{u}^+(n)\|_1 + \|\hat{u}^-(n)\|_1 \;\le\; h^d \big(\boldsymbol{|}\hat{\mathbf{u}}_n^+\boldsymbol{|}_{R1} \,+\, \boldsymbol{|}\hat{\mathbf{u}}_n^-\boldsymbol{|}_{R1}\big).$$

Proof: Let us consider the mapping $\Phi_n(R) : l(G_n(R)) \mapsto E_n(R, \mathbb{R}^d)$ defined by $u(n) = \sum_{\boldsymbol{k}} u_{\boldsymbol{k}} \psi_{\boldsymbol{k}}$. In terms of functions $\chi_{\boldsymbol{k}i+}$ of Subsection 3.2 we easily get:

$$\partial_i \, u(n) \;=\; \sum_{\boldsymbol{k} \in I_n(R)} \big(U_i(p_i) \mathbf{u}_n\big)_{\boldsymbol{k}} \, \chi_{\boldsymbol{k}i+}. \tag{26}$$

Apparently we have for $p = 1$ and $p = \infty$ the following inequalities: $\|u\|_p \le h^{d/p} \boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{Rp}$, $\|\partial_i u\|_p \le h^{d/p} \boldsymbol{|}U_i(p_i) \mathbf{u}_n\boldsymbol{|}_{Rp}$. Hence, by the Riesz-Thorrin theorem we get (i).

Similarly we prove (ii). For $p = 1$ and $p = \infty$ we have $h^{d/p} \boldsymbol{|}\hat{\mathbf{u}}_n\boldsymbol{|}_{Rp} \le \|u\|_p$. Now we use the expression $\psi_{\boldsymbol{k}+p\boldsymbol{e}_i}(\boldsymbol{x}) = \psi_{\boldsymbol{k}}(\boldsymbol{x} - ph\boldsymbol{e}_i)$ and get

$$\big(U_i(p) \hat{\mathbf{u}}_n\big)_{\boldsymbol{k}} \;=\; \frac{1}{\|\psi_{\boldsymbol{k}}\|_1} \, (\psi_{\boldsymbol{k}} \,|\, \eth_i(p) \, u).$$

Because of $\|\, \eth_i(p) \, u\|_p \le \|\partial_i u\|_p$ for $p = 1, \infty$, we have $h^{d/p} \boldsymbol{|}U_i(p) \hat{\mathbf{u}}_n\boldsymbol{|}_{Rp} \le \|\partial_i u\|_p$. Assertion (ii) follows now from the Riesz-Thorrin theorem.

The right hand side of double inequality (iii) is implied by (i) while the left hand side is implied by (ii). Therefore we have to prove the equality $\|u\|_1 = \|u^+(n)\|_1 + \|u^-(n)\|_1$. This equality is a consequence of $\sum_{\boldsymbol{k}} \psi_{\boldsymbol{k}} = 1$ on $\mathbb{R}^d$ and $\|\widehat{u^+}(n)\|_1 = \sum_{\boldsymbol{k}} (\psi_{\boldsymbol{k}} | u^+)$. **QED**

4 Construction of discretizations

It is important to underline at the beginning that discretizations A_n of differential operator $A(\boldsymbol{x})$ are defined prior to discretizations of the forms (5), (6). This fact is in a full agreement with construction of discretizations A_n in this section. Some classes of discretizations are derived from a general principle which is not based on finite difference formulas and cannot be apriori related to variational equalities. Nevertheless, bilinear forms must be associated to A_n so that A_n are derived from the corresponding variational equalities. The constructed bilinear forms are considered as discretizations of the original form (3). These forms are basic objects in our proof of convergence of approximate solutions. In the next two subsections schemes and the corresponding discretized forms are constructed for two classes of methods.

Forms $a_n(\cdot, \cdot)$ on $E(R, \mathbb{R}^d) \times E(R, \mathbb{R}^d)$ and matrices A_n on $G_n(R)$ are related by equalities:

$$a_n(v, u) \;=\; \langle \, \mathbf{v} \,|\, A_n \, \mathbf{u} \, \rangle_R.$$

In addition, the discretized forms determine the discretized variational equalities:

$$\lambda \langle \, \mathbf{v}_n \,|\, \mathbf{u}_n \, \rangle_R \;+\; a_n(v, u) \;=\; \langle \, \mathbf{v}_n \,|\, \boldsymbol{\mu}_n \, \rangle_R, \quad \mathbf{v}_n \;\in\; w_{2,1}(G_n(R)). \tag{27}$$

To discretize $A(\boldsymbol{x})$ means to associate to $A(\boldsymbol{x})$ a sequence of matrices A_n on $G_n(R), n \in \mathbb{N}$. Of course, the matrices A_n must be constructed reasonably in order to enable demonstrations of the convergence of numerical solutions. The convergence analysis is postponed until two next sections. Therefore, in this section, the terminology "discretizations" of $A(\boldsymbol{x})$ instead of approximations of $A(\boldsymbol{x})$ seams to be more suitable.

Discretizations to be considered in this section are possible if certain conditions on a_{ij} are fulfilled. The required conditions are stronger than in Assumption 2.1. By relaxing them gradually as $n \to \infty$ we obtain discretizations for a general $A(\boldsymbol{x})$ given by Assumption 2.1.

To a given diffusion tensor $a = \{a_{ij}\}_{11}^{dd}$ we associate an auxiliary tensor $\hat{a}$ defined by the expressions:

$$\hat{a}_{ii} \;=\; a_{ii}, \quad \hat{a}_{ij} \;=\; -|a_{ij}| \quad i \ne j. \tag{28}$$

Assumption 4.1 (on discretization conditions)

1. There exist $\boldsymbol{q} \in \mathbb{N}^d$*, a finite index set* $\mathfrak{L}$ *and a partition* $\mathbb{R}^d = \cup_l D_l$*, where the sets* D_l *are connected unions of* $C_n(\boldsymbol{q}, \boldsymbol{x}), \boldsymbol{x} \in G_n(Q)$*. The sets* D_l *and the diffusion tensor* $a = \{a_{ij}\}_{11}^{dd}$ *must be mutually related as follows:*

- *The tensor-valued function* $\boldsymbol{x} \mapsto a(\boldsymbol{x})$ *is continuous on* D_l *and the functions* $a_{ij}, i \neq j$ *do not change signs on* D_l*.*
- *There is a grid-step* $h_\varepsilon = 2^{-n(\varepsilon)}$ *and the closed sets*

$$D_l(\varepsilon) \;=\; \cup_{\boldsymbol{x} \in cls(D_l)} S_{n(\varepsilon)}(\boldsymbol{q}, \boldsymbol{x}),$$

such that the functions a_{ij} *can be extended to* $D_l(\varepsilon)$*, not changing the signs on* $D_l(\varepsilon)$*, and the strict ellipticity (2) is valid on* $D_l(\varepsilon)$ *with the same bounds* $\underline{M}, \overline{M}$*.*

2. For each $l \in \mathfrak{L}$ *the auxiliary diffusion tensor* $\hat{a}$ *is strictly positive definite on* $D_l(\varepsilon)$*.*

3. To each D_l *there is associated parameter* $\boldsymbol{p}(l) \in \mathbb{N}^d$*, such that the following inequality is valid:*

$$\begin{aligned}\omega(a) \;=\; & \inf_n \min_l {}_i \inf \Big\{ \frac{1}{p_i(l)} \inf_{\boldsymbol{z} \in S_n(\boldsymbol{p}(l), \boldsymbol{x})} a_{ii}(\boldsymbol{z}) \\ & - \sum_{m \neq i} \frac{1}{p_m(l)} \sup_{\boldsymbol{z} \in S_n(\boldsymbol{p}(l), \boldsymbol{x})} |a_{im}(\boldsymbol{z})| \;:\; \boldsymbol{x} \in G_n \cap D_l(\varepsilon) \Big\} \;>0.\end{aligned}$$

Condition 3. is crucial in our construction of discretizations A_n which have a particular feature called the compartmental structure. In the next definition I is an index set, and matrices $A = \{a_{ij}\}_{II}$ are considered in linear spaces $l_p(I)$ consisting of functions on the set I:

DEFINITION 4.1 (Compartmental structure) *A matrix* A *in* $l_\infty(I)$ *is said to be of positive type if* $A = pI - B$, $p > 0$, $B \geq 0$ *and* $\|B\|_\infty \leq p$*. It is called conservative if* $B\mathbf{1} = p\mathbf{1}$*. A matrix* A *in* $l_1(I)$ *is said to have the compartmental structure if* $A = pI - B$, $B \geq 0$ *and* $\|B\|_1 \leq p$*. It is called conservative if for each* $\mathbf{u} \geq \mathbf{0}$ *there holds* $\|B\mathbf{u}\|_1 = p\|\mathbf{u}\|_1$*.*

LEMMA **4.1** *Let* A *in* $l_1(I)$ *be compartmental. Then*

(i) If A *is conservative then* $\|B^m\|_1 = p^m$ *for each* $m \in \mathbb{N}$*.*

(ii) The spectrum $sp(A)$ *of a compartmental matrix is contained in* $\Re\lambda \geq 0$*. If* A *is conservative, then* $0 \in sp(A)$*.*

In the case of $d = 2$ Conditions 2. and 3. can be always fulfilled. For $d > 2$ there exist positive definite diffusion tensors $\{a_{ij}\}_{11}^{dd}$ such that Conditions 2. and 3. are not possible [LR3].

Discretizations A_n are defined in terms of its matrix entries $(A_n)_{\boldsymbol{kl}}$, where $h\boldsymbol{k}, h\boldsymbol{l} \in G_n$. For a fixed $\boldsymbol{x} = h\boldsymbol{k} \in G_n$ the set of all the grid-knots $\boldsymbol{y} = h\boldsymbol{l}$ such that $(A_n)_{\boldsymbol{kl}} \neq 0$ is denoted by $\mathcal{N}(\boldsymbol{x})$ and called the *numerical neighbourhood* of A_n at $\boldsymbol{x} \in G_n$. In our constructions the sets $\mathcal{N}(\boldsymbol{x})$ for $\boldsymbol{x} \in G_n \cap D_l$ are mutually alike. A set $\mathcal{N}(\boldsymbol{x})$ contains always a "cross" consisting of $\boldsymbol{x}$ and $2d$ elements $\pm hp_i(l)\boldsymbol{e}_i$. Additional elements of $\mathcal{N}(\boldsymbol{x})$ depend on the sign of $a_{ij}, i \neq j$.

Two classes of discretizations are analyzed. One of these classes can be defined straightforwardly in terms of forward and backward difference formulas. The resulting discretizations are called *basic schemes.* Discretizations of the other class come from a general principle [LR3] and they are called *extended schemes.*

The compartmental structure of discretizations A_n of the differential operator $A(\boldsymbol{x})$ is the goal of overall analysis. Here we describe a general approach to the constructions of discretizations A_n with the compartmental structure which is based on reduction to the two-dimensional problems.

Figure 1: Asumption on function a_{12}



$a_{12} = 0$ on $\cup C_n(\boldsymbol{p}, \boldsymbol{v})$
$p_1 = 2,\ p_2 = 1$

The index set of pairs $I(d) = \{\{ij\} : i < j, i, j = 1, 2, \ldots, d, i \neq j\}$ has the cardinal number $m(d) = d(d-1)/2$. To each index $\{kl\} \in I(d)$ we associate three coefficients,

$$a_{kk}^{\{kl\}} = \frac{1}{d-1} a_{kk}, \quad a_{ll}^{\{kl\}} = \frac{1}{d-1} a_{ll}, \quad a_{kl}^{\{kl\}} = a_{kl}, \tag{29}$$

and a bilinear form $a^{\{kl\}}(\cdot, \cdot)$,

$$a^{\{kl\}}(v, u) \;=\; \sum_{i,j \in \{r,s\}} \int_D a_{ij}^{\{kl\}}(\boldsymbol{x})\, \partial_i v(\boldsymbol{x})\, \partial_j u(\boldsymbol{x})\, d\boldsymbol{x}. \tag{30}$$

Apparently, for each pair $v, u \in C^{(1)}(\mathbb{R}^d)$ with compact supports, the following equality is valid:

$$a(v, u) \;=\; \sum_{\{kl\} \in I(d)} a^{\{kl\}}(v, u).$$

To each of the forms $a^{\{kl\}}(\cdot, \cdot)$ we must associate a sequence of forms $a_n^{\{kl\}}(\cdot, \cdot)$ and matrices $A_n^{\{kl\}}$ constructed by two-dimensional schemes. Then the matrix

$$A_n \;=\; \sum_{\{kl\} \in I} A_n^{\{kl\}}, \tag{31}$$

is a discretization of $A_0(\boldsymbol{x})$. If each $A_n^{\{kl\}}$ has the compartmental structure then A_n also has the compartmental structure. However, A_n can have the compartmental structure although no $A_n^{\{kl\}}$ is compartmental. Condition 3. of Assumption 4.1 ensures this advantageous property in our constructions.

4.1 Two methods of discretizations

The forms must be constructed by the rules (29)-(30) having in mind that the construction for higher dimensional cases depends on the construction for two-dimensional case. Therefore, we are due to describe the construction for the two-dimensional case.

Basic schemes

In the case of numerical grids $G_n(P)$ the numerical schemes and corresponding discrete bilinear forms can be easily mutually related. A proof of convergence in Sobolev spaces for various right hand sides are presented in our works [LR2, LR3]. In the present case we extend analysis to problems with the function a_{12} having both signs. Difficulties appear at those grid-knots where the sign of a_{12} changes.

The discretizations are constructed by assuming that the sets $\{\boldsymbol{x} \in \mathbb{R}^d : a_{12}(\boldsymbol{x}) < 0\}$ and $\{\boldsymbol{x} \in \mathbb{R}^d : a_{12}(\boldsymbol{x}) > 0\}$ are separated by a set (connected) so that it contains a connected subset which is equal to the union of cubes $C_n(\boldsymbol{p}, \boldsymbol{x}), \boldsymbol{x} \in G_n(P)$ for some $n \in \mathbb{N}$. An illustration of this assumption is given in Figure 1. This assumption is not valid for a general matrix-valued function $\boldsymbol{x} \mapsto a(\boldsymbol{x})$. Therefore we have to comprehend this assumption as a step of an approximation procedure in our process of construction of discretizations A_n. For the sake of simple and brief presentation we assume in the next construction that the assumption is valid already for $n = 1$.

Let us define the sets $D_n(-) \subset \mathbb{R}^d$ by:

$$D_n(-) \; = \; \{\cup_{\boldsymbol{v} \in G_n(P)} C_n(\boldsymbol{p}, \boldsymbol{v}) \; : \; a_{12}(\boldsymbol{x}) \leq 0 \text{ for } \boldsymbol{x} \in C_n(\boldsymbol{p}, \boldsymbol{v})\},$$

and $D_n(+) = D_n(-)^c$. Now we define the subgrids $G_n(P, -)$ consisting of all the vertices $\boldsymbol{v} \in G_n(P)$ which determine the set $D_n(-)$. As well we need the subgrid $G_n(P, +)$ consisting of those grid-knots $\boldsymbol{x}$ for which the segments $I(\boldsymbol{x}, \boldsymbol{y}), \boldsymbol{y} = \boldsymbol{x} + h_n p_1 \boldsymbol{e}_1$ have the following property $I(\boldsymbol{x}, \boldsymbol{y}) \subset cls(D_n(+))$. It is easy to verify that each $\boldsymbol{x} \in G_n(P)$ must be contained in one of sets $G_n(P, \mp)$ and each of segments $I_i(\boldsymbol{x}, \boldsymbol{y}), \boldsymbol{y} = \boldsymbol{x} + h p_i \boldsymbol{e}_i$ must have both of its end points $\boldsymbol{x}, \boldsymbol{y}$ in some $G_n(P, \mp)$. Some of grid-knots and some of segments are contained in both sets, $G_n(P, \mp)$ and $cls(D_n(\mp))$, respectively. For a segment in $\boldsymbol{e}_2$-direction, $I(\boldsymbol{x}_1, \boldsymbol{x}_2), \boldsymbol{x}_2 = \boldsymbol{x}_1 + h_n p_2 \boldsymbol{e}_2$, the following is true. If $\boldsymbol{x}_1 \in G_n(P, -)$ then $I(\boldsymbol{x}_1, \boldsymbol{x}_2) \subset cls(D_n(-))$. If $\boldsymbol{x}_2 \in G_n(P, +)$ then $I(\boldsymbol{x}_2, \boldsymbol{x}_1)$ (downward vertical segment) may be outside of $cls(D_n(+)))$. If this happens then this segment is contained in a cube $cls(C_n(\boldsymbol{p}, \boldsymbol{z}))$ on which the function a_{12} has zero values. This fact is a consequence of a strict separation of supports of functions $\max\{a_{12}, 0\}$ and $\min\{a_{12}, 0\}$ and will be utilized in our constructions of discrete bilinear forms.

To define forms and entries we need

$$\begin{array}{rcl} \boldsymbol{z}^{(\pm)} & = & p_i \boldsymbol{e}_i \, \pm \, p_j \boldsymbol{e}_j \; \in I_n, \\ \boldsymbol{x}^{(\pm +)}(n) & = & \frac{1}{2} \big(\pm h \, p_i \boldsymbol{e}_i \, + \, h \, p_j \boldsymbol{e}_j \big) \; \in S_n(\boldsymbol{p}, \boldsymbol{0}), \\ \boldsymbol{x}^{(\pm -)}(n) & = & \frac{1}{2} \big(\pm h \, p_i \boldsymbol{e}_i \, - \, h \, p_j \boldsymbol{e}_j \big) \; \in S_n(\boldsymbol{p}, \boldsymbol{0}). \end{array}$$

Obviously we have $h\boldsymbol{z}^{(\alpha\beta)} \in G_n$ while $\boldsymbol{x}^{(\pm)}(n)$ are not necessary in G_n.

Let us consider a sequence of two-dimensional forms on $E_n(R, \mathbb{R}^2) \times E_n(R, \mathbb{R}^2)$ which are defined by the following expressions:

$$\begin{array}{rcl} a_n(v, u) & = & a_n^{(-)}(v, u) \, + \, a_n^{(+)}(v, u) \\ a_n^{(-)}(v, u) & = & \sum_{i\,j=1}^2 \sum_{\boldsymbol{x} \in G_n(P,-)} \big(\partial_i(p_i h) v \big)(\boldsymbol{x}) \\ & \times & a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(++)}(n)) \, \big(\partial_j(p_j h) u \big)(\boldsymbol{x}), \\ a_n^{(+)}(v, u) & = & \sum_{i\,j=1}^2 \sum_{\boldsymbol{x} \in G_n(P,+)} \big(\partial_i((-1)^{i-1} p_i h) v \big)(\boldsymbol{x}) \\ & \times & a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(+-)}(n)) \, \big(\partial_j((-1)^{j-1} p_j h) u \big)(\boldsymbol{x}). \end{array} \tag{32}$$

Discretizations A_n of differential operator $A_0(\boldsymbol{x})$ can be easily obtained from the constructed forms variationaly. Let us define matrices $\mathcal{A}_n^{(\pm)}(i, j, p_i, p_j)$ as the diagonal matrices with entries $\mathcal{A}_n^{(\pm)}(i, j, p_i, p_j)_{\boldsymbol{x}\boldsymbol{x}} = a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(+\pm)}(n))$. In terms of matrices $U_i(p_i), V_i(p_i)$ and $Z_n(p_i, i)$ of (14) we get the following expressions:

$$\begin{array}{rcl} A_n & = & - \displaystyle\sum_{l \in \mathcal{L}_-} \sum_{ij=1}^{2} V_i(p_i) \, \mathcal{A}_n^{(+)}(i, j, p_i, p_j) \, U_j(p_j) \, \mathbb{1}_{G_n(P,-)} \\ & & - \displaystyle\sum_{l \in \mathcal{L}_+} \sum_{ij=1}^{2} V_i(p_i) \, \Lambda_i^T \, \mathcal{A}_n^{(-)}(i, j, p_i, p_j) \, \Lambda_j \, U_j(p_j) \, \mathbb{1}_{G_n(P,+)}, \end{array} \tag{33}$$

where $\Lambda_1 = I, \Lambda_2 = Z_n(-p_2, 2)$ and $\mathbb{1}_{G_n(P,\mp)}$ are the projectors on the linear subspaces of grid-functions with supports in the sets $G_n(P, \mp)$, respectively. The entries of A_n can be also easily calculated for grid-knots of $int(G_n(P, \mp))$. In order to get simple expressions we use the following abbreviations:

$$a_{ij}^{(\alpha\beta)} \; = \; a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(\alpha\beta)}(n)), \quad \alpha, \beta \in \{+, -\}.$$

Figure 2: Numerical neighbourhoods at two grid-knots of $G_n(P,\mp)$



The entries on the cross $\boldsymbol{x} \pm h_r p_r \boldsymbol{e}_r, r = 1, 2$ have the structure:

$$\begin{aligned}
\left(A_n\right)_{\boldsymbol{kk}\pm p_1\boldsymbol{e}_1} &= -\frac{1}{h^2p_1^2}\begin{cases} a_{11}^{(\pm+)} - \dfrac{p_1}{p_2}\left|a_{12}^{(\pm+)}\right| & \text{for} \quad a_{12} \leq 0, \\ a_{11}^{(\pm-)} - \dfrac{p_1}{p_2}\left|a_{12}^{(\pm-)}\right| & \text{for} \quad a_{12} \geq 0, \end{cases} \\
\left(A_n\right)_{\boldsymbol{kk}\pm p_2\boldsymbol{e}_2} &= -\frac{1}{h^2p_2^2}\left[a_{22}^{(+\pm)} - \frac{p_2}{p_1}\left|a_{12}^{(+\pm)}\right|\right].
\end{aligned}$$

The entries in the plane spanned by $\boldsymbol{e}_1, \boldsymbol{e}_2$ have the structure:

$$\begin{aligned}
\left(A_n\right)_{\boldsymbol{kk}\pm\boldsymbol{z}^{(-)}} &= -\frac{1}{h^2p_1p_2}\left|a_{12}^{(\pm\mp)}\right| \quad \text{for} \quad a_{12} \leq 0 \text{ on } D_l, \\
\left(A_n\right)_{\boldsymbol{kk}\pm\boldsymbol{z}^{(+)}} &= -\frac{1}{h^2p_1p_2}\left|a_{12}^{(\pm\pm)}\right| \quad \text{for} \quad a_{12} \geq 0 \text{ on } D_l,
\end{aligned}$$

where $+\boldsymbol{z}^{(\pm)}$ is associated with the upper and $-\boldsymbol{z}^{(\pm)}$ with the lower indices of $a^{(\alpha\beta)}$. The diagonal entries $(A_n)_{\boldsymbol{kk}}$ are equal to the negative sum of all the entries $(A_n)_{\boldsymbol{kl}}$, $\boldsymbol{l} \neq \boldsymbol{k}$. Entries for grid-knots at $bnd(G_n(P,\mp))$ can be more complex. In two-dimensional problems the forms (32) are natural and the entries of A_n calculated from variational equalities (27) can be perceived as buildups made by forward/backward finite difference formulæ. This approach is self-understanding and we call it the standard approach. The result are basic schemes of discretization. An illustration of numerical neighbourhoods of basic schemes is given in Figure 2. These schemes are studied thoroughly in [SMMM]. The strict ellipticity of forms $a_n(\cdot,\cdot)$ is expressed always in terms of the original pair of positive numbers $\underline{M}, \overline{M}$. Generally, the strict ellipticity of discretized forms follow from the compartmental structure of A_n. These problems are analyzed in the next section.

Extended schemes

Contrary to the standard approach in developing discretizations of $A_0(\boldsymbol{x})$ we have methods based on some general principle and which are not a priori related to the forward/backward finite difference formulas. One of such methods is described here. Principles of construction are given in [LR3]. To each D_l we must associate elements $\boldsymbol{r}(l) \in \mathbb{N}$ playing the role analogous to $\boldsymbol{p}$ for basic schemes.

Again we assume the strict separation of sets $F(\pm) = \operatorname{supp}(\max\{\pm a_{12}, 0\})$. The set $\mathfrak{L}$ of Assumption 4.1 is partitioned into the subsets $\mathfrak{L}_\mp$, where $l \in \mathfrak{L}_-$ means that $a_{ij} \leq 0$ on D_l and $l \in \mathfrak{L}_+$ means $a_{ij} \geq 0$ on D_l. Let us remind that the sets D_l in present case are determined in terms of cubes $C_{n(\varepsilon)}(\boldsymbol{r}(l), \boldsymbol{x})$. There is always a room of arbitrariness in a determination of these sets. The following maximal property removes some of arbitrariness. There exist a $n(\varepsilon) \in \mathbb{N}$ such that the sets $D_l, l \in \mathfrak{L}_-$ have the following properties:

a) $a_{12} \leq 0$ on $\cup_{l\in\mathfrak{L}_-} D_l$,

b) Each D_l satisfies Assumption 4.1,

c) Each $C_{n(\varepsilon)}(\boldsymbol{r}(l),\boldsymbol{x}), l \in \mathfrak{L}_-, \boldsymbol{x} \in G_n$ on which $a_{12} = 0$ is contained in some of $D_l, l \in \mathfrak{L}_-$.

Now we can define $D(-) = \cup_{l\in\mathfrak{L}_-} D_l$ and $D(+) = D(-)^c$. In our proceeding discussion we assume $n(\varepsilon) = 1$. The subgrids $G_n(-) = G_n \cap D(-)$ have the same properties as the corresponding subrids $G_n(P,-), p_1 = p_2 = 1$ in the subsection on basic schemes. Similarly, the sets $G_n(+) = G_n \cap D(+)$ coincide with $G_n(P,+)$ as well. Now we define subgrids $G_n(l,-) = G_n(-) \cap D_l$ and conclude that $G_n(l,-)$ form a partition of $G_n(-)$. However, the sets $G_n(l,+) = G_n(+) \cap D_l$ do not form a partition of $G_n(+)$ because some of grid-knots at $\partial D(+)$ may be outside of each $G_n(+,l)$. Therefore, we have to extend the sets D_l to wider sets $\tilde{D}_l$ such that $G_n(l,+) = G_n(+) \cap \tilde{D}_l$ form a partition of $G_n(+)$. The sets $\tilde{D}_l$ cannot be defined uniquely. Here we demand the following properties. The sets must be disjoint, and the closure of $int(\tilde{D}_l)$ must coincide with $cls(D_l)$. Thus we have:

$$G_n(l,-) \;=\; G_n(-) \cap D_l, \quad G_n(l,+) \;=\; G_n(+) \cap \tilde{D}_l.$$

In accordance with our discussion about properties of sets $G_n(P,\pm)$ in the subsection on basic schemes we finally conclude that $G_n(l,\pm)$ cover G_n and some of them may have common grid-knots.

The forms $a_n(\cdot,\cdot)$ are expressed in terms of $a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(\pm\pm)}(l,n)), \boldsymbol{x} \in G_n$, where $\boldsymbol{x}^{(\pm\pm)}(l,n)$ are certain elements in $\mathbb{R}^d$. Since $\boldsymbol{x}^{(\pm\pm)}(l,n)$ can take values outside of $cls(D_l)$ we are due to specify how to take values of $a_{ij}(\boldsymbol{x} + \boldsymbol{x}^{(\pm\pm)}(l,n))$ in such cases. The values must be taken in the set $D_l(\varepsilon)$ of Assumption 4.1. In this way we conclude that the entries of A_n are calculated in terms of values of coefficients a_{ij} at points which are not grid-knots.

It is convenient to use a representation $a_n(v,u) = a_n^{(-)}(v,u) + a_n^{(+)}(v,u)$, where the forms $a_n^{(\mp)}(v,u)$ are related to the index sets $\mathfrak{L}_\mp$ as previously. Let us define

$$\begin{aligned}
\boldsymbol{t}^{(\pm+)}(\boldsymbol{r}) \;&=\; \tfrac{1}{2}\big(\pm h\, r_i(l)\boldsymbol{e}_i \,+\, h\, r_j(l)\boldsymbol{e}_j\big) \;\in S_n(\boldsymbol{r},\boldsymbol{0}),\\
\boldsymbol{t}^{(\pm-)}(\boldsymbol{r}) \;&=\; \tfrac{1}{2}\big(\pm h\, r_i(l)\boldsymbol{e}_i \,-\, h\, r_j(l)\boldsymbol{e}_j\big) \;\in S_n(\boldsymbol{r},\boldsymbol{0}).
\end{aligned}$$

Obviously $\boldsymbol{x}^{(\alpha\beta)}(l,n)$ and $\boldsymbol{t}^{(\alpha\beta)}(\boldsymbol{r})$ coincide for $\boldsymbol{p} = \boldsymbol{r}(l)$. The form $a_n^{(-)}(\cdot,\cdot)$ is defined by:

$$\begin{aligned}
a_n^{(-)}(v,u) \;=\; &\sum_{l\in\mathfrak{L}_-}\ \sum_{\boldsymbol{x}\in G_n(l)} \Big(\sum_{i=1}^{2} a_{ii}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{1}))\,\big(\daleth_i(h)v\big)(\boldsymbol{x})\big(\daleth_i(h)u\big)(\boldsymbol{x})\\
&+ \sum_{i\neq j} a_{ij}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r}))\,\big(\daleth_i(r_i(l)h)v\big)(\boldsymbol{x})\big(\daleth_j(r_j(l)h)u\big)(\boldsymbol{x})\\
&+ \sum_{i\neq j} a_{ij}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r}))\,\frac{r_i(l)h}{r_j(l)h}\,\Big[\big(\daleth_i(h)v\big)(\boldsymbol{x})\big(\daleth_i(h)u\big)(\boldsymbol{x})\\
&- \big(\daleth_i(r_i(l)h)v\big)(\boldsymbol{x})\big(\daleth_i(r_i(l)h)u\big)(\boldsymbol{x})\Big]\,\Big).
\end{aligned} \tag{34}$$

We obtain $a_n^{(+)}(v,u)$ from $a_n^{(-)}(v,u)$ by replacing $\daleth_i(r_i(l)h)$ with $\daleth_i((-1)^{i-1}(r_i(l)h))$, $\mathfrak{L}_-$ with $\mathfrak{L}_+$ and $\boldsymbol{t}^{(++)}$ with $\boldsymbol{t}^{(+-)}$. The forms $a_n^{(\mp)}(v,u)$ are not second degree polynomials of $\daleth_i(h)$ with simple structure. Due to the compartmental structure of A_n they can be ultimately represented as forms depending on $\daleth_i(q_i h)$ with various q_i. For the quantities $a_n^{(\mp)}(u,u)$ more comprehensible expressions can be written down such as (47) and (48).

Discretizations A_n have a general expression:

$$A_n \;=\; \sum_{l\in\mathfrak{L}_-} A_n^{(-)}(l)\,\mathbb{1}_{G_n(l,-)} \;+\; \sum_{l\in\mathfrak{L}_+} A_n^{(+)}(l)\,\mathbb{1}_{G_n(l,+)},$$

where

$$
\begin{aligned}
A_n^{(-)}(l) = &\; -\sum_{i=1}^{2} V_i\,\mathcal{A}_n^{(+)}(i,i,1,1)\,U_i \;-\; \sum_{i\neq j}^{2} V_i(r_i(l))\,\mathcal{A}_n^{(+)}(i,j,r_i,r_j)\,U_j(r_j(l)) \\
& -\sum_{i\neq j}^{2} \frac{r_i(l)h}{r_j(l)h}\Big[V_i\,\mathcal{A}_n^{(+)}(i,j,1,1)\,U_i \;-\; V_i(r_i(l))\,\mathcal{A}_n^{(+)}(i,j,r_i,r_j)\,U_i(r_i(l))\Big], \\
A_n^{(+)}(l) = &\; -\textstyle\sum_{i=1}^{2} V_i\,\Lambda_i(r_i)^T\,\mathcal{A}_n^{(-)}(i,i,1,1)\,\Lambda_i(r_i)\,U_i \\
& -\displaystyle\sum_{i\neq j}^{2} V_i(r_i(l))\,\Lambda_i(r_i)^T\mathcal{A}_n^{(-)}(i,j,r_i,r_j)\Lambda_j(r_j)\,U_j(r_j(l)) - \sum_{i\neq j}^{2}\frac{r_i(l)h}{r_j(l)h} \\
& \times\Big[V_i\,\Lambda_i(r_i)^T\,\mathcal{A}_n^{(-)}(i,j,1,1)\,\Lambda_i(r_i)\,U_i \;-\; V_i(r_i(l))\,\mathcal{A}_n^{(-)}(i,j,r_i,r_j)\,U_i(r_i(l))\Big],
\end{aligned}
\tag{35}
$$

and where $\Lambda_1(r_1) = I, \Lambda_2(r_2) = Z_n(-r_2(l), 2)$ as in the previous case.

In order to write down the entries of A_n we need the following abbreviations:

$$
\begin{aligned}
\boldsymbol{w}^{(\pm)}(l) &= r_i(l)\boldsymbol{e}_i \pm r_j(l)\boldsymbol{e}_j \;\in I_n, \\
a_{ij}^{(\alpha\beta)}(\boldsymbol{r}) &= a_{ij}(\boldsymbol{x} + \boldsymbol{t}^{(\alpha\beta)}(\boldsymbol{r})), \quad \alpha,\beta \in \{+,-\}, \\
\hat{a}_{12}^{(-+)}(\boldsymbol{r}) &= a_{12}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r}) - h\boldsymbol{e}_1), \\
\hat{a}_{12}^{(+-)}(\boldsymbol{r}) &= a_{12}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r}) - h\boldsymbol{e}_2), \\
\hat{a}_{ii}^{(++)}(\boldsymbol{r}) &= a_{ii}^{(++)}(\boldsymbol{r}), \quad \hat{a}_{ii}^{(--)}(\boldsymbol{r}) \;=\; a_{ii}^{(--)}(\boldsymbol{r}).
\end{aligned}
$$

Then we have the following nontrivial off-diagonal entries of A_n:

$$
\begin{aligned}
\big(A_n\big)_{\boldsymbol{kk}\pm\boldsymbol{e}_1} &= -\frac{1}{h^2}\begin{cases} a_{11}^{(\pm+)}(\boldsymbol{1}) - \dfrac{r_1(l)}{r_2(l)}\big|\hat{a}_{12}^{(\pm+)}(\boldsymbol{r})\big| & \text{for} \quad a_{12} \le 0, \\ a_{11}^{(\pm-)}(\boldsymbol{1}) - \dfrac{r_1(l)}{r_2(l)}\big|\hat{a}_{12}^{(\pm-)}(\boldsymbol{r})\big| & \text{for} \quad a_{12} \ge 0, \end{cases} \\
\big(A_n\big)_{\boldsymbol{kk}\pm\boldsymbol{e}_2} &= -\frac{1}{h^2}\Big[a_{22}^{(+\pm)}(\boldsymbol{1}) - \frac{r_2(l)}{r_1(l)}\big|\hat{a}_{12}^{(+\pm)}(\boldsymbol{r})\big|\Big].
\end{aligned}
\tag{36}
$$

The entries in the plane spanned by $\boldsymbol{e}_1, \boldsymbol{e}_2$ have the structure:

$$
\begin{aligned}
\big(A_n\big)_{\boldsymbol{kk}\pm\boldsymbol{w}^{(-)}(l)} &= -\frac{1}{h^2 r_1(l) r_2(l)}\big|a_{12}^{(\pm\mp)}(\boldsymbol{r})\big| \quad \text{for} \quad a_{12} \le 0 \text{ on } D_l, \\
\big(A_n\big)_{\boldsymbol{kk}\pm\boldsymbol{w}^{(+)}(l)} &= -\frac{1}{h^2 r_1(l) r_2(l)}\big|a_{12}^{(\pm\pm)}(\boldsymbol{r})\big| \quad \text{for} \quad a_{12} \ge 0 \text{ on } D_l,
\end{aligned}
\tag{37}
$$

where $+\boldsymbol{w}^{(\pm)}$ is associated with the upper and $-\boldsymbol{w}^{(\pm)}$ with the lower indices of $a^{(\alpha\beta)}$, respectively.

Discretizations A_n defined by (36), (37) are called extended schemes. The numerical neighbourhoods are illustrated in Figure 3.

In (35) we have 4 sums with respect to the indices i, j. The first and second sums have expressions similar to Expressions (32). They contribute to Expressions (37) and to a part of entries in (36). Unfortunately there appear non-trivial entries $(A_n)_{\boldsymbol{kl}}, \boldsymbol{l} = \boldsymbol{k} \pm r_i\boldsymbol{e}_i$. These entries must be canceled by contributions from the third and fourth sums. These two latter sums lack the structure similar to (32) since the sum includes the terms $a_{12}\reflectbox{$\partial$}_1 v \reflectbox{$\partial$}_1 u$ and $a_{12}\reflectbox{$\partial$}_2 v \reflectbox{$\partial$}_2 u$. So the net result of all four sums are entries (37). Let $J \subset I_n(l)$ consists of grid-knots in some of $\boldsymbol{x} \in G_n(l,\pm)$ for which the numerical neighbourhoods $\mathcal{N}(\boldsymbol{x})$ have only internal grid-knots, $\mathcal{N}(\boldsymbol{x}) \subset int(G_n(l,\pm))$. The associated diagonal submatrix $(A_n)_{JJ}$ is symmetric. Diagonal submatrices for which $\mathcal{N}(\boldsymbol{x})$ contain boundary grid-knots of $G_n(l,\pm)$ may lack the symmetry. If the quantities $\hat{a}_{ij}^{(\alpha\beta)}$ in (36) are replaced with $a_{ij}^{(\alpha\beta)}$, the symmetry of A_n is lost altogether, although the convergence is still preserved. However, the quantities $a_{ij}^{(\alpha\beta)}$ must not be replaced with $a_{ij}(h\boldsymbol{k})$ since the resulting $(A_n)_{\boldsymbol{kl}}$ would be discretizations of $-\sum_{ij} a_{ij}\partial_i\partial_j$.

Figure 3: Numerical neighbourhoods at internal grid-knots of D_l



$a_{12} \geq 0$ $a_{12} \leq 0$

$r_1 = 3,\ r_2 = 1$

4.2 Discretization with compartmental structure

Now we can describe general structure of constructed discretizations of $A_0(\boldsymbol{x})$. From the definition of bilinear forms $a_n(u,v) = \langle \mathbf{v} | A_n \mathbf{u} \rangle$ the following property is obvious: If $\mathbf{v} = \mathbf{1}$ and $\mathbf{u}$ are with compact supports on G_n there must be $a_n(1,u) = a_n(u,1) = 0$, implying that the row sums and column sums of A_n have zero values. Hence, if the off-diagonal entries of A_n are non-positive then the matrices A_n are simultaneously M-matrices and have the compartmental structure. We consider here only the extended scheme.

It is convenient to utilize the quantities:

$$\omega_n(a_{ii}, \boldsymbol{x}) \;=\; \frac{1}{d-1} \sum_{s \neq i}^{d} a_{ii}(h\boldsymbol{k} + h\boldsymbol{m}_{ii}(l,s)), \tag{38}$$

where $\boldsymbol{m}_{ii}(l,s)$ are defined by the rules of construction of extended schemes.

Discretization procedure 4.1 *Let Assumption 4.1 be valid and matrices A_n on G_n be constructed by the rule (31). Then their entries have the following properties:*

1. *Entries of $(A_n)_{\boldsymbol{kl}}, \boldsymbol{k}, \boldsymbol{l} \in I_n$, $\boldsymbol{x} = h\boldsymbol{k}$ are linear combinations of $a_{ij}(\boldsymbol{x}_{ij}(n,\boldsymbol{x},l))$ where $\boldsymbol{x}_{ij}(n,\boldsymbol{x},l) = h\boldsymbol{k} + h\boldsymbol{m}_{ij}(l,s)$, $\boldsymbol{m}_{ij}(l,s)$ are n-independent elements of $\mathbb{R}^d$ and $l \in \mathfrak{L}, s \in \{1,2,\ldots,d\}$.*

2. *For each grid-knot $\boldsymbol{x} = h\boldsymbol{k}$: $(A_n)_{\boldsymbol{kk}} = -\sum_{\boldsymbol{l}} (A_n)_{\boldsymbol{kl}}$.*

3. *For each $\boldsymbol{x} = h\boldsymbol{k} \in cls(D_l)$ entries on the "cross branches" $\boldsymbol{x} \pm h\boldsymbol{e}_i$, i.e. $(A_n)_{\boldsymbol{k} \pm h\boldsymbol{e}_i}$, are defined by:*

$$\begin{aligned}(A_n)_{\boldsymbol{xx} \pm h\boldsymbol{e}_i} \;&=\; -\frac{1}{h^2}\Big[\omega_n(a_{ii},\boldsymbol{x}) \\ &\quad - \sum_{m \neq i} \frac{r_i(l)}{r_m(l)} \left| a_{im}(\boldsymbol{x}_{im}(n,\boldsymbol{x},l)) \right| \Big].\end{aligned}$$

4. *Entries of A_n which are not on the "cross branches" are defined by using elements $\boldsymbol{z}_{ij}(l) = r_i(l)\boldsymbol{e}_i - r_j(l)\boldsymbol{e}_j \in I_n$ or elements $\boldsymbol{z}_{ij}(l) = r_i(l)\boldsymbol{e}_i + r_j(l)\boldsymbol{e}_j \in I_n$:*

$$(A_n)_{\boldsymbol{kk} \pm \boldsymbol{z}_{ij}(l)} \;=\; -\frac{1}{r_i(l) r_j(l)} \left| a_{ij}(\boldsymbol{x}_{ij}(n,\boldsymbol{x},l)) \right|.$$

Some peculiar features regarding the structure of sets $\mathcal{N}(\boldsymbol{x}), \boldsymbol{x} = h\boldsymbol{k} \in G_n \cap D_l$, must be pointed out. If $a_{ij}, i \neq j$ is not changing the sign in a neighbourhood of $\boldsymbol{x}$ then the minimal number of elements in $\mathcal{N}(\boldsymbol{x})$ is $1 + d + d^2$. In this case the set $\mathcal{N}(\boldsymbol{x})$ consists of its center, $2d$-grid-knots on the d-dimensional cross $\{\pm \boldsymbol{e}_i : i = 1,2,\ldots,d\}$ and 2 grid-knots in each two-dimensional plane. Generally, the number of grid-knots in two-dimensional plane may be larger than 2. Here we consider only the case of two grid-knots at most in the two-dimensional planes. This demand has the following implication on the construction of discretizations $A_n^{(rs)}$. Let the pairs $\boldsymbol{e}_r, \boldsymbol{e}_s$ and $\boldsymbol{e}_s, \boldsymbol{e}_t$

define two-dimensional planes and let $A_n^{(rs)}, A_n^{(st)}$ be the corresponding discretizations which are constructed using the parameters $\boldsymbol{r}^{(rs)}, \boldsymbol{r}^{(st)}$. Then there must hold $(\boldsymbol{r}^{(rs)})_s = (\boldsymbol{r}^{(st)})_s$. In such case the off-diagonal entries of A_n have the structure as described in 4. of Discretization procedure 4.1. The described structure of sets $\mathcal{N}(\boldsymbol{x})$ is valid for all $\boldsymbol{x}$ because the functions $a_{ij}, i \neq j$ do not change sign on $S_n(\boldsymbol{p}, \boldsymbol{x})$. This is an important consequence of the strict separation of sets $\max\{a_{12}, 0\}$.

Obviously that all the constructed forms a_n of this section are discretizations of the form (3). One can be easily convinced that the terminology "a discretization of the original form (3)" is not artificial. At the present level of analysis it is easy to check $a(v,u) = \lim_n h^d a_n(v,u)$ for any pair $v, u \in C_0^{(1)}(\mathbb{R}^d)$.

THEOREM **4.1** *Let Assumption 4.1 be valid. There exist discretizations A_n which are constructed by the rules of Discretization procedure 4.1, such that A_n have the compartmental structure.*

PROOF: For each D_l we have to choose the parameters $r_i(l)$ of the properties 3. and 4. of Discretization procedure 4.1 so that the condition 3. of Assumption 4.1 is valid. The uniform continuity of coefficients on D_l and the condition 3. of Assumption 4.1 ensure the compartmental structure of matrices A_n as demonstrated in [LR3]. **QED**

5 Convergence in W_2^1-spaces

Discretizations of the original variational problem (5) or (6) are defined in terms of a sequence of bilinear forms $a_n(\cdot,\cdot)$ on $E_n(R,\mathbb{R}^d) \times E_n(R,\mathbb{R}^d)$ and a sequence of linear functionals $\langle \cdot | \boldsymbol{\mu}_n \rangle_R$ on $E_n(R,\mathbb{R}^d)$. The associated discretized variational problems are defined by equalities (27). The discretized variational equalities (27) can be rewritten in an equivalent manner:

$$(\lambda I + A_n)\,\mathbf{u}_n = \boldsymbol{\mu}_n, \tag{39}$$

where in the case of Problem (6) $\mathbf{u}_n, \boldsymbol{\mu}_n$ are grid-functions on G_n and in the case of Problem (5) they are grid-functions on $G_n(R,D)$ or $G_n(R,D)$.

Problems (5) or (6) are solved numerically in two steps. In the first step we construct grid-functions $\mathbf{u}_n$ on $G_n(R)$ or $G_n(R,D)$ according to (39). The obtained grid-functions represent the solution at grid-knots, therefore, we call them *grid-solutions*. The grid-solutions are imbedded into the spaces $E_n(R,\mathbb{R}^d)$ or $E_n(R,D)$ by (16), and the convergence $u(n) \to u$ must be proved in some Banach spaces. Therefore we call functions $u(n)$ *approximate solutions*.

Though the functions $u(n) = \Phi(R)\mathbf{u}_n$ are called approximate solutions, this terminology has to be justified after a convergence analysis. The convergence proofs are based on some properties of the forms $a_n(\cdot,\cdot)$ and linear functionals $\langle \cdot | \boldsymbol{\mu}_n \rangle_R$ to be described in details later in this section. Most of the analysis in this section is related to Problem (6) on $\mathbb{R}^d$. The obtained results can be easily applied to Problem (5) on a bounded domain. This is carried out at the end of section.

5.1 Consistency

Certain number of notions important for the convergence of approximate solutions is formulated in terms of sequences of functions with a particular structure:

$$\begin{aligned} \mathfrak{V} &= \{v(n) \,:\, n \in \mathbb{N}\} \subset \cup_n E_n(R,\mathbb{R}^d), \\ \mathfrak{U} &= \{u(n) \,:\, n \in \mathbb{N}\} \subset \cup_n E_n(R,\mathbb{R}^d). \end{aligned} \tag{40}$$

DEFINITION 5.1 (Consistency) *The forms $a_n(\cdot,\cdot)$ on $E_n(R,\mathbb{R}^d) \times E_n(R,\mathbb{R}^d)$ are consistent with the form (3) if*

$$a(v,u) = \lim_n h^d\, a_n(v(n), u(n))$$

is valid for any pair $\mathfrak{V}, \mathfrak{U}$ of (40) such that $\mathfrak{V}$ converges weakly in $W_2^1(\mathbb{R}^d)$ to v, and $\mathfrak{U}$ converges strongly in $W_2^1(\mathbb{R}^d)$ to u.

PROPOSITION **5.1** *Let a sequence of matrices $\{A_n : n \in \mathbb{N}\}$ be constructed by basic schemes or extended schemes and let $\{a_n(\cdot,\cdot) : n \in \mathbb{N}\}$ be the corresponding sequence of discretizations of (3). Then the forms a_n are consistent with the form (3).*

This important result is proved by a lemma which is formulated bellow. Let $\mathfrak{V}$ and $\mathfrak{U}$ be defined by (40) and converge weakly and strongly in $W_2^1(\mathbb{R}^d)$ to $v, u \in W_2^1(\mathbb{R}^d)$, respectively. Expressions $\omega_n(a_{ij}, \boldsymbol{x})$ for $i = j$ are defined by (38). In this proof we extend this definition to the case $i \neq j$ and define $\omega_n(a_{ij}, \boldsymbol{x}) = a_{ij}(\boldsymbol{x}_{ij}(n, \boldsymbol{x}, l))$, where $\boldsymbol{x}_{ij}(n, \boldsymbol{x}, l) = \boldsymbol{x} + h\, \boldsymbol{m}_{ij}(l, n, s)$ are constructed by the rules of basic and extended schemes. For each $l \in \mathfrak{L}$ and each pair $i, j \in \{1, 2, \ldots, d\}$ there holds:

$$\lim_n \int_{D_l(\varepsilon)} a_{ij}(\boldsymbol{x})\, (\partial_i v(n))(\boldsymbol{x})\, (\partial_j u(n))(\boldsymbol{x})\, d\boldsymbol{x}$$
$$= \lim_n \sum_{\boldsymbol{x} \in G_n(P) \cap D_l(\varepsilon)} \omega_n(a_{ij}, \boldsymbol{x}) \int_{C_n(\boldsymbol{p}(l), \boldsymbol{x})} (\partial_i v(n))(\boldsymbol{x})\, (\partial_j u(n))(\boldsymbol{x})\, d\boldsymbol{x},$$

Let us point out that the written identity is unchanged if we replace $\omega_n(a_{ij}, \boldsymbol{x})$ with $a_{ij}(\boldsymbol{x})$. Let us consider a bilinear form on $W_2^1(\mathbb{R}^d) \times W_2^1(\mathbb{R}^d)$ defined by

$$\theta_n(l, v, u) \;=\; \sum_{\boldsymbol{x} \in G_n(R) \cap D_l} p(\boldsymbol{w}(n, \boldsymbol{x})) \int_{C_n(\boldsymbol{r}, \boldsymbol{x})} \partial_i v(\boldsymbol{y})\, \partial_j u(\boldsymbol{y}) d\boldsymbol{y}, \tag{41}$$

where p is a uniformly continuous function on D_l and $\boldsymbol{w}(n, \cdot)$ is a transformation of $\mathbb{R}^d$ such that $\boldsymbol{w}(n, \boldsymbol{x}) \in C_n(\boldsymbol{r}, \boldsymbol{x})$. Then

$$\lim_n \theta_n(l, v(n), u(n)) \;=\; \int_{D_l} p(\boldsymbol{y})\, \partial_i v(\boldsymbol{y})\, \partial_j u(\boldsymbol{y}) d\boldsymbol{y} \tag{42}$$

for any pair $\mathfrak{V}, \mathfrak{U}$ of (40), converging in $W_2^1(\mathbb{R}^d)$ weakly to v and strongly to u, respectively.

The object of next analysis is the bilinear functional on $E_n(R, \mathbb{R}^d) \times E_n(R, \mathbb{R}^d)$ defined by:

$$\gamma_n(l, v, u) \;=\; h^d \operatorname{vol}(R) \sum_{\boldsymbol{x} \in G_n(R) \cap D_l} p(\boldsymbol{w}(n, \boldsymbol{x}))\, \beth_i(r_i h) v(h\boldsymbol{k})\, \beth_j(r_j h) u(h\boldsymbol{k}). \tag{43}$$

In particular:

$$\gamma_n(l, v(n), u(n)) \;=\; h^d \operatorname{vol}(R) \sum_{\boldsymbol{x} \in G_n(R) \cap D_l} p(\boldsymbol{w}(n, \boldsymbol{x}))\, v_{\boldsymbol{k}i}\, u_{\boldsymbol{k}j},$$

where $v_{\boldsymbol{k}i} = \beth_i(r_i h) v(h\boldsymbol{k}), u_{\boldsymbol{k}j} = \beth_j(r_j h) u(h\boldsymbol{k})$.

LEMMA **5.1** *Let the sequence $\mathfrak{V}$ converge weakly in $W_2^1(\mathbb{R}^d)$ to v and $\mathfrak{U}$ converge strongly to u in $W_2^1(\mathbb{R}^d)$. Then*

$$\lim_n \gamma_n(l, v(n), u(n)) \;=\; \lim_n \theta_n(l, v(n), u(n)) \;=\; \int_{D_l} p(\boldsymbol{x})\, \partial_i v(\boldsymbol{x})\, \partial_j u(\boldsymbol{x})\, d\boldsymbol{x}.$$

PROOF: We have to analyze $\theta_n(l, v(n), u(n))$ as $n \to \infty$. To simplify notation we assume $i, j \in \{1, 2\}$. The indices $\boldsymbol{k} = (k_1, k_2, \ldots, k_s)$ are denoted shortly as $\boldsymbol{k} = (k_1, \boldsymbol{k}')$ and $\boldsymbol{k} = (k_1, k_2, \boldsymbol{k}'')$, $k_1, k_2 \in \mathbb{Z}$. The corresponding r_1, r_2 are denoted by r, s, respectively. After inserting expressions for $v(R, n), u(R, n)$ into (41) and carrying out a straightforward calculation we get expressions:

$$\theta_n(l, v(n), u(n)) \;=\; \sum_{\boldsymbol{x} \in G_n(R) \cap D_l} p(\boldsymbol{w}(n, \boldsymbol{x}))\, \rho_n(h\boldsymbol{k}, v(n), u(n)),$$

where $\rho_n(h\boldsymbol{k}, v(n), u(n))$ denotes an integral which for the case of $i = j = 1$ has the form:

$$\begin{aligned}\rho_n(h\boldsymbol{k}, v(n), u(n)) \;&=\; \int_{C_n(\boldsymbol{r},h\boldsymbol{k})} \partial_1 v(R,n)(\boldsymbol{y})\,\partial_1 u(R,n)(\boldsymbol{y})\,d\boldsymbol{y} \;=\; \sum_{\boldsymbol{k}'\boldsymbol{l}'} (\psi_{\boldsymbol{k}'}|\psi_{\boldsymbol{l}'}) \\ &\times \left[\sum_{l=k,k+r} v_{k\boldsymbol{k}'} u_{l\boldsymbol{l}'}\,(\partial_1\psi_k|\partial_1\psi_l) \;+\; \sum_{l=k,k+r} v_{k+r\boldsymbol{k}'} u_{l\boldsymbol{l}'}\,(\partial_1\psi_{k+r}|\partial_1\psi_l)\right] \\ &=\; rh \sum_{\boldsymbol{k}'\boldsymbol{l}'} (\psi_{\boldsymbol{k}'}|\psi_{\boldsymbol{l}'})\,\big(\eth_1(rh)v\big)(h\boldsymbol{k})\,\big(\eth_1(rh)u\big)(h\boldsymbol{k}).\end{aligned}$$

Now we use quantities $s_{\boldsymbol{kl}} = (\psi_{\boldsymbol{k}}|\psi_{\boldsymbol{l}})\|\psi_{\boldsymbol{k}}\|_1^{-1}$ and bring into mind their properties $s_{\boldsymbol{kl}} \geq 0, \sum_{\boldsymbol{l}} s_{\boldsymbol{kl}} = 1$.

$$\theta_n(l, v(n), u(n)) = h^d\,\mathrm{vol}(R) \sum_{k,\boldsymbol{k}',\boldsymbol{l}'\in I_n(R),\boldsymbol{x}\in D_l} p(\boldsymbol{w}(n,\boldsymbol{x}))\,s_{\boldsymbol{k}'\boldsymbol{l}'}\,v_{(k\boldsymbol{k}')i}\,u_{(k\boldsymbol{l}')i}.$$

Analogously we get for $i = 1, j = 2$:

$$\begin{aligned}\theta_n(l, v(n), u(n)) \;=\; & h^{d-2}\,\mathrm{vol}(R'') \sum_{k,l,\boldsymbol{k}'',\boldsymbol{l}''\in I_n(R),\boldsymbol{x}\in D_l} s_{\boldsymbol{k}''\boldsymbol{l}''}\,p(\boldsymbol{w}(n,h\boldsymbol{k})) \int_{J(k,r)\times J(l,s)} dz_1 dz_2 \\ &\times \Big[v_{(k,l,\boldsymbol{k}'')i}\,u_{(k,l,\boldsymbol{l}'')j}\,\psi_k(z_1)\psi_l(z_2) \;+\; v_{(k,l,\boldsymbol{k}'')i}\,u_{(k+r,l,\boldsymbol{l}'')j}\,\psi_{k+r}(z_1)\psi_l(z_2) \\ &+ v_{(k,l+s,\boldsymbol{k}'')i}\,u_{(k,l,\boldsymbol{l}'')j}\,\psi_k(z_1)\psi_{l+s}(z_2) \;+\; v_{(k,l+s,\boldsymbol{k}'')i}\,u_{(k+r,l,\boldsymbol{l}'')j}\,\psi_{k+r}(z_1)\psi_{l+s}(z_2)\Big],\end{aligned}$$

where $J(k,r) = [hk, h(k+r)]$ and R'' stands for the parameter set $\{\boldsymbol{r}_0, r_3, r_4, \ldots, r_d\}$. Upon integration over z_1, z_2 we get

$$\theta_n(l, v(n), u(n)) = \frac{1}{4}\,h^{d-2}\,\mathrm{vol}(R'') \sum_{k,l,\boldsymbol{k}'',\boldsymbol{l}''\in I_n(R),\boldsymbol{x}\in D_l} s_{\boldsymbol{k}''\boldsymbol{l}''} p(\boldsymbol{w}(n,h\boldsymbol{k}))$$
$$\Big[v_{(k,l,\boldsymbol{k}'')i}\,u_{(k,l,\boldsymbol{l}'')j} + v_{(k,l,\boldsymbol{k}'')i} u_{(k+r,l,\boldsymbol{l}'')j} + v_{(k,l+s,\boldsymbol{k}'')i} u_{(k,l,\boldsymbol{l}'')j} + v_{(k,l+s,\boldsymbol{k}'')i} u_{(k+r,l,\boldsymbol{l}'')j}\Big].$$

We finish the proof for the case $i = j = 1$ since the case $i \neq j$ can be treated analogously. The quantity γ_n would be equal to θ_n if $s_{\boldsymbol{kl}}$ were absent and the double sum were replaced with the single sum over indices $\boldsymbol{k}$. Therefore we are due to estimate their difference:

$$\gamma_n(l, v(n), u(n)) - \theta_n(l, v(n), u(n)) \;=\; h^d\,\mathrm{vol}(R) \sum_{\boldsymbol{kl}} p(\boldsymbol{w}(n,h\boldsymbol{k}))\,v_{\boldsymbol{k}i}\,\delta_{kl}\,s_{\boldsymbol{k}'\boldsymbol{l}'}\,\big(u_{\boldsymbol{l}j} - u_{\boldsymbol{k}j}\big). \tag{44}$$

To estimate the right hand side we need $\overline{p} = \sup p$:

$$\begin{gathered}\Big|\gamma_n(l, v(n), u(n)) \;-\; \theta_n(l, v(n), u(n))\Big| \;\leq \\ \overline{p}\,h^{d/2}\big\| U_i(r_i)\mathbf{v}_n \big\|_{R2}\,h^{d/2}\,\max\{\big\|\big(Z(r_j,j) - I\big)U_j(r_j)\mathbf{u}_n\big\|_{R2} \,:\, j = 1, 2, \ldots, d\}.\end{gathered}$$

By Theorem 3.1 we have

$$\begin{aligned} h^d\,\big\|U_i(r_i)\mathbf{v}_n\big\|_{R2}^2 \;&\leq\; (1-\sigma^2)^{-1}\,\|\,\partial_i\,v(R,n)\|_2^2, \\ h^d\,\big\|(Z(r_j,j) - I)\,U_j(r_j)\mathbf{u}_n\big\|_{R2}^2 \;&\leq\; (1-\sigma^2)^{-1}\,\|(Z(\boldsymbol{w}) - I)\,\partial_j\,u(R,n)\|_2^2,\end{aligned}$$

where $\boldsymbol{w} = hr_j\boldsymbol{e}_j$. Due to the strong convergence of $\mathfrak{U}$ we have

$$\begin{aligned} h^{d/2}\,\big\|\big(Z(r_j,j) - I\big)U_j(r_j)\mathbf{u}_n\big\|_{R2} \;&\leq\; (1-\sigma^2)^{-1/2}\,\|\big(Z(\boldsymbol{w}) - I\big)\,\partial_j\,u(n,R)\|_2 \\ &\leq\; (1-\sigma^2)^{-1/2}\,\Big[\|(Z(\boldsymbol{w}) - I)\,\partial_j\,u\|_2 \,+\, 2\,\|\,\partial_j\,u - \,\partial_j\,u(R,n)\|_2\Big] \;\to\; 0,\end{aligned}$$

so that $\lim_n \gamma_n(l, v(n), u(n)) = \lim_n \theta_n(l, v(n), u(n)) = \int p\partial_i v\partial_j u$. **QED**

Now a proof of Proposition 5.1 follows from the inequalities

$$\lim_n \sum_{l\in\mathfrak{L}'} \gamma_n(l, v(n), u(n)) \;=\; \int_{\cup\{D_l : l\in\mathfrak{L}'\}} a_{ij}(\boldsymbol{x})\,\partial_i v(\boldsymbol{x})\partial_j u(\boldsymbol{x})\,d\boldsymbol{x}.$$

in which $p(\boldsymbol{w}(n,h\boldsymbol{k}))$ of (43) is replaced with $a_{ij}(n,\boldsymbol{x},l)$.

5.2 Strict ellipticity of discretized forms

A discrete form $a_n(\cdot,\cdot)$ on $l_0(G_n(R)) \times l_0(G_n(R))$ is said to be strictly elliptic [Yo] if there exist two positive numbers $\underline{M}(a_n), \overline{M}(a_n)$ such that

$$\underline{M}(a_n) \sum_{i=1}^{d} \| U_i(r_i)\mathbf{u} \|_{R2}^2 \;\leq\; a_n(u,u) \;\leq\; \overline{M}(a_n) \sum_{i=1}^{d} \| U_i(r_i)\mathbf{u} \|_{R2}^2 .$$

For a sequence of discrete forms $a_n(\cdot,\cdot)$ we need a stronger result. The strict ellipticity must be uniform with respect to n and $\mathrm{vol}(R) = \prod r_i$ different values of $\boldsymbol{r}_0$ in the parameter set $R = (\boldsymbol{r}_0, \boldsymbol{r})$.

DEFINITION 5.2 *Let A_n be discretized by the rules of Discretization procedure 4.1. Discrete forms $a_n(\cdot,\cdot)$ on $l_0(G_n(R)) \times l_0(G_n(R))$ are said to be strictly elliptic uniformly with respect to $n \in \mathbb{N}$ if there exist positive numbers $\underline{M} \leq \overline{M}$ such that*

$$\underline{M} \sum_{i=1}^{d} \| U_i(r_i)\mathbf{u} \|_{R2}^2 \;\leq\; a_n(\mathbf{u}_n,\mathbf{u}_n)_R \;\leq\; \overline{M} \sum_{i=1}^{d} \| U_i(r_i)\mathbf{u} \|_{R2}^2 . \tag{45}$$

for all $n \in \mathbb{N}$ and all $0 \leq (\boldsymbol{r}_0)_i < r_i, i = 1,2,\ldots,d$.

PROPOSITION **5.2** *Let the discretizations A_n of $A_0(\boldsymbol{x}) = -\sum \partial_i a_{ij}(\boldsymbol{x})\partial_j$ be constructed by the rules of Discretization procedure 4.1. If A_n have the compartmental structure then the discrete forms $\mathbf{v},\mathbf{u} \mapsto \langle \mathbf{v} | A_n \mathbf{u} \rangle_R$ are strictly elliptic on $l_0(G_n(R)) \times l_0(G_n(R))$ uniformly with respect to $n \in \mathbb{N}$.*

PROOF: First we consider the basic scheme for a two-dimensional grid. From the structure of bilinear forms (32) we have

$$a_n(u,u) \;\geq\; \underline{M} \| U_1(p_1)\mathbf{u}_n \|_2^2 + \underline{M} \Big(\| K(-)U_2(-p_2)\mathbf{u}_n \|_2^2 + \| K(+)\Lambda_2 U_2(-p_2)\mathbf{u}_n \|_2^2 \Big), \tag{46}$$

where $K(\mp) = \mathbb{1}_{G_n(P,\mp)}$ are projectors. Let us consider a pair $\boldsymbol{x}, \boldsymbol{y} = \boldsymbol{x} - hp_2\boldsymbol{e}_2$ which is involved in the definition of operator $U_2(-p_2)$. The indices of $\boldsymbol{x},\boldsymbol{y}$ are $\boldsymbol{k},\boldsymbol{l}$, respectively. If $\boldsymbol{x},\boldsymbol{y} \in G_n(P,+)$ then $(U_2(-p_2)\mathbf{u}_n)_{\boldsymbol{k}}$ can be replaced with $(U_2(p_2)\mathbf{u}_n)_{\boldsymbol{l}}$. If $\boldsymbol{y} \notin G_n(P,+)$ then $\boldsymbol{y} \in G_n(P,-)$. In this case the term $(U_2(-p_2)\mathbf{u}_n)_{\boldsymbol{l}}$ can be omitted from the sum in (46) because this term is already contained in the corresponding sum of $U_1(p_1)\mathbf{u}_n$. Hence, the right hand side of (46) can be estimated from bellow by $\underline{M}\sum_{i=1}^{2} \| U_i(p_i)\mathbf{u}_n \|_2^2$. In this way the left hand side of inequality (45) is proved. The right hand side of this double inequality follows by choosing the double value of $\overline{M}$. In the case of $d > 2$ we use the construction (31) and get the same lower and upper bounds. Let us point out that the compartmental structure is not used in this step of proof.

Let us now consider a two-dimensional problem with an extended scheme. In the present step, the compartmental structure is utilized in an essential way. The form $a_n^{(-)}(u,u)$ of (34) can be rewritten as:

$$\begin{aligned} &a_n^{(-)}(u,u) \;= \\ &\sum_{l \in \mathcal{L}_-} \sum_{\boldsymbol{k} \in G_n(l,-)} \sum_{i \neq j} \Big(a_{ii}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\mathbf{1})) + \frac{r_i(l)}{r_j(l)} a_{12}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r})) \Big) (\exists_i(h)u)(\boldsymbol{x})^2 \;- \\ &\sum_{l \in \mathcal{L}_-} \sum_{\boldsymbol{k} \in G_n(l,-)} a_{12}(\boldsymbol{x} + \boldsymbol{t}^{(++)}(\boldsymbol{r})) \left(\sqrt{\frac{r_1(l)}{r_2(l)}} (\exists_1(r_1h)u)(\boldsymbol{x}) - \sqrt{\frac{r_2(l)}{r_1(l)}} (\exists_2(r_2h)u)(\boldsymbol{x}) \right)^2 . \end{aligned} \tag{47}$$

Let us remind that $a_{12} \leq 0$ on the set $D(-)$. Due to the compartmental structure the first term is positive definite. The second term is positive semidefinite and can be disregarded in the next step of estimation from bellow. The result is:

$$a_n^{(-)}(u,u) \;\geq\; \omega(a) \sum_{l \in \mathcal{L}_-} \sum_{\boldsymbol{k} \in G_n(l,-)} \sum_{i} (U_i \mathbf{u}_n)_{\boldsymbol{k}}^d .$$

where $\omega(a)$ is the positive number specified in Assumption 4.1. For the form $a_n^{(+)}(u,u)$ we have an analogous inequality involving the summation over all the indices $\boldsymbol{k} \in G_n(+)$.

$$a_n^{(+)}(u,u) =$$
$$\sum_{l\in\mathcal{L}_+}\sum_{\boldsymbol{k}\in G_n(l,+)}\sum_{i\neq j}\Big(a_{ii}(\boldsymbol{x}+\boldsymbol{t}^{(+-)}(\mathbf{1}))-\frac{r_i(l)}{r_j(l)}a_{12}(\boldsymbol{x}+\boldsymbol{t}^{(+-)}(\boldsymbol{r}))\Big)(\partial_i((-1)^{i-1}h)u)(\boldsymbol{x})^2+$$
$$\sum_{l\in\mathcal{L}_+}\sum_{\boldsymbol{k}\in G_n(l,+)} a_{12}(\boldsymbol{x}+\boldsymbol{t}^{(+-)}(\boldsymbol{r}))\left(\sqrt{\frac{r_1(l)}{r_2(l)}}(\partial_1(r_1h)u)(\boldsymbol{x})-\sqrt{\frac{r_2(l)}{r_1(l)}}(\partial_2(-r_2h)u)(\boldsymbol{x})\right)^2. \tag{48}$$

The lower bound follows in the same way as for $a_n^{(-)}$. **QED**

5.3 W_2^1-convergence

We have shown how the form $\lambda(v|u)+a(v,u)$ is discretized by forms $h^d\lambda\langle\,\mathbf{v}_n|\mathbf{u}_n\,\rangle_R + \langle\,\mathbf{v}_n|A_n\mathbf{u}_n\,\rangle_R$. In order to solve discretized problem (39), we have to describe a discretization of the linear function $v \to \langle\,v|\mu\,\rangle$ by $h^d\langle\,\mathbf{v}_n|\boldsymbol{\mu}_n\,\rangle_R$, where $\boldsymbol{\mu}_n \in l(G_n(R))$. First we must demonstrate the existence of $\boldsymbol{\mu}_n$ such that $h^d\langle\,\mathbf{v}_n|\boldsymbol{\mu}_n\,\rangle_R \to \langle\,v|\mu\,\rangle$. Discretizations of μ are defined by (22) so that (24) is valid.

LEMMA **5.2** *Let μ be a continuous linear functional on $W_2^1(\mathbb{R}^d)$. There exists discretizations $\boldsymbol{\mu}_n(R) \in l(G_n(R))$ such that*

$$\langle\,u(n)\,|\,\mu\,\rangle \;=\; h^d\,\langle\,\mathbf{u}_n\,|\,\boldsymbol{\mu}_n\,\rangle_R \;=\; h^d\,\mathrm{vol}(R)\sum_{\boldsymbol{k}\in I_n(R)} u_{\boldsymbol{k}}\mu_{\boldsymbol{k}}. \tag{49}$$

Obviously, for each sequence $\{u(n) : n \in \mathbb{N}\}$ weakly converging to some $u \in W_2^1(\mathbb{R}^d)$ the following equality holds: $\lim_n \langle\,u(n)|\mu\,\rangle = \langle\,u|\mu\,\rangle$.

From this Lemma we have

$$|\langle\,\mathbf{u}_n|\boldsymbol{\mu}_n\,\rangle_R| \;\le\; h^{-d/2}\,q_R(\mathbf{u}_n)^{1/2}\,\|\mu\|_{2,-1}.$$

Inequalities (45), (49) and the variational equalities (27) imply the first result towards our proof of convergence of approximate solutions. If $\mathbf{u}_n$ solve (27) or (39) then

$$h^d\,\underline{M}\,\boldsymbol{|}\mathbf{u}_n\boldsymbol{|}_{2,1}^2 \;\le\; h^d\,\langle\,\mathbf{u}_n\,|\,(\lambda I + A_n)\,\mathbf{u}_n\,\rangle_R \;\le\; \|u(R,n)\|_{2,1}\,\|\mu\|_{2,-1}. \tag{50}$$

COROLLARY **5.1** *Let $\mathbf{u}_n = T(\lambda, A_n)\boldsymbol{\mu}_n$ and $u(R,n) = \Phi_n(R)\mathbf{u}_n$. Then for each R the sequence $\mathfrak{U} = \{u(R,n) : n \in \mathbb{N}\} \subset \cup_n E_n(R,\mathbb{R}^d)$ converges weakly in $W_2^1(\mathbb{R}^d)$ to some $u \in W_2^1(\mathbb{R}^d)$.*

Let u^* be the solution of (6). Then the sequence of functions $\hat{u}^*(n)$, defined by (20), strongly converges to u^* in $W_2^1(\mathbb{R}^d)$. In the remaining part of this analysis we have to demonstrate the expected property $\lim_n u(R,n) = \lim_n \hat{u}^*(R,n) = u^*$ for each R. We follow the well-known finite element technique.

$$\begin{array}{rcl} \underline{M}\,h^d\,\boldsymbol{|}\mathbf{u}_n-\hat{\mathbf{u}}_n^*\boldsymbol{|}_{R2,1}^2 & \le & h^d\,\langle\,\mathbf{u}_n-\hat{\mathbf{u}}_n^*\,|\,(\lambda I + A_n)\,(\mathbf{u}_n-\hat{\mathbf{u}}_n^*)\,\rangle_R \\ & = & h^d\,\langle\,\mathbf{u}_n-\hat{\mathbf{u}}_n^*\,|\,(\lambda I + A_n)\,\mathbf{u}_n\,\rangle_R \\ & - & h^d\,\langle\,\mathbf{u}_n-\hat{\mathbf{u}}_n^*\,|\,(\lambda I + A_n)\,\hat{\mathbf{u}}_n^*\,\rangle_R \\ = h^d\;\langle\,\mathbf{u}_n-\hat{\mathbf{u}}_n^*\,|\,\boldsymbol{\mu}_n\,\rangle_R & - & h^d\,\langle\,\mathbf{u}_n-\hat{\mathbf{u}}_n^*\,|\,(\lambda I + A_n)\,\hat{\mathbf{u}}_n^*\,\rangle_R. \end{array} \tag{51}$$

By Lemma 5.2 the first term on the right hand side converges to $\langle\,u-u^*|\mu\,\rangle$. By the consistency property of Proposition 5.1 the second term converges to the same value.

THEOREM **5.1** *Let $\mathfrak{U}$ be as in Corollary 5.1. Then the sequence $\mathfrak{U}$ converges $W_2^1(\mathbb{R})$-strongly to the unique solution u^* to (6).*

From this result, Lemma 5.2 and Lemma 3.1 we get another important result for $\lambda = 0$.

COROLLARY **5.2** *Let D be a bounded domain with Lipsithz boundary and $\mu \in W_2^{-1}(D)$. Let $A_n(D)$ be the restriction to $G_n(R, D)$ of A_n, $\boldsymbol{\mu}_n$ on $G_n(R, D)$ satisfy (49) and $\mathbf{u}_n = A_n(D)^{-1}\boldsymbol{\mu}_n$. Then the sequence $\mathfrak{U}$ converges strongly in $\dot{W}_2^1(D)$ to the unique weak solution u of (5).*

6 Convergence in L_1-spaces

In this section we consider Problem (5) for a bounded domain D and numerical solutions in $\dot{W}_p^1(D)$. Furthermore, for the sake of simplicity of exposition, we restrict our analysis to the differential operator $A_0(\boldsymbol{x}) = -\sum_{ij} \partial_i a_{ij}(\boldsymbol{x}) \partial_j$ and its discretizations $A_n(D)$. We consider the boundary value problem (5) with $\mu \in \mathcal{R}(D)$ and its discretizations

$$A_n(D)\,\mathbf{u}_n \;=\; \boldsymbol{\mu}_n \tag{52}$$

on $G_n(R, D)$, where $A_n(D)$ have the compartmental structure. The discretizations $\boldsymbol{\mu}_n \in l(G_n(R, D))$ are defined by

$$\mu_{\boldsymbol{k}} \;=\; \frac{\psi_{\boldsymbol{k}}}{\|\psi_{\boldsymbol{k}}\|_1}\,\langle\, \psi_{\boldsymbol{k}} \,|\, \mu\,\rangle.$$

Apparently, $\langle\, v(n)|\mu\,\rangle = h^d \langle\, \mathbf{v}_n|\boldsymbol{\mu}_n\,\rangle_R$ for any $v(n) = \sum v_{\boldsymbol{k}}\psi_{\boldsymbol{k}} \in E_n(R, D)$. Problem (52) is defined for grid-functions on $G_n(R, D)$.

6.1 Boundedness of approximate solutions

For any pair $r, R,\ 0 < r < R \le 1$ and $\boldsymbol{v} \in D$ we define the balls $B_r(\boldsymbol{v}), B_R(\boldsymbol{v})$. By using the functions $u \in \dot{W}_2^1(D)$ we define the sets $A(r, s, \boldsymbol{v}) = \{\boldsymbol{x} \in B_r(\boldsymbol{v}) \,:\, u(\boldsymbol{x}) \ge s\}$, where $s \in \mathbb{R}$. The measure of $A(r, s, \boldsymbol{v})$ is denoted by $a(r, s, \boldsymbol{v}) = \operatorname{meas}(A(r, s, \boldsymbol{v}))$. Let us assume that there exist a subset $\mathcal{G} \subset \dot{W}_2^1(D)$ and two numbers, c_1, c_2, independent of $r, R, s, \boldsymbol{v}$, such that the inequality

$$\sum_{i=1}^{d} \|\, \mathbb{1}_{A(r,s,\boldsymbol{v})}\, \partial_i u \,\|_2^2 \;\le\; c_1 \|\, \mathbb{1}_{A(R,s,\boldsymbol{v})}\, (u{-}s)\,\|_2^2 \left[1 \;+\; \frac{1}{(R-r)^2}\right] {+} c_2 s^2 \big(a(R, s, \boldsymbol{v})\big)^{1-2/q} \tag{53}$$

is valid for a fixed $q > d$, each $u \in \mathcal{G}$, and all $s \ge s_0$ with some s_0. Then [LU] the functions $u \in \mathcal{G}$ are bounded on D with a bound which depends on D, c_1, c_2 and $\|u\|_2$. Instead of balls $B_r(\boldsymbol{v})$ one can use rectangles $S_n(\boldsymbol{p}, \boldsymbol{v}) = \prod_{i=1}^{d}[-hp_i + hv_i, hv_i + hp_i]$ as well. Actually, it is sufficient to consider balls (rectangles) with centers $\boldsymbol{v}$ in a dense set of D whose radii (edges) are contained in a sequence $\{r_m : m \in \mathbb{N}\} \subset (0, 1)$, converging to zero. For instance, the sets $S_n(\boldsymbol{p}, \boldsymbol{v})$ and $\boldsymbol{v} \in G_n(R)$, match this weaker condition. This fact enables a straightforward application of (53) to the sequence of functions in Corollary 5.2. First, we have to establish a discretized version of (53), and then we have to prove that the constructed discretized version implies (53) for the sequence $\mathfrak{U}$ of Corollary 5.2. Further, let

$$\begin{aligned} A(\boldsymbol{r}, s, \boldsymbol{v}) \;&=\; S_m(\boldsymbol{p}, \boldsymbol{v}) \,\cap\, \{\, u(\boldsymbol{x}) \,\ge\, s \,\}, \\ A(\boldsymbol{R}, s, \boldsymbol{v}) \;&=\; S_{m-t}(\boldsymbol{p}, \boldsymbol{v}) \,\cap\, \{\, u(\boldsymbol{x}) \,\ge\, s \,\}, \end{aligned}$$

where the sets $A(\boldsymbol{R}, s, \boldsymbol{v})$ are larger than $A(\boldsymbol{r}, s, \boldsymbol{v})$, $\operatorname{dist}(\partial A(\boldsymbol{r}, s, \boldsymbol{v}), \partial A(\boldsymbol{R}, s, \boldsymbol{v})) = h(m)(2^t - 1)\underline{p}$, where $\underline{p} = \min p_k$ and $m > t$. Hence, the symbols $\boldsymbol{r}, \boldsymbol{R}$ stand for the d-dimensional parameters $h(m)\boldsymbol{p}$ and $h(m-t)\boldsymbol{p}$, respectively. The discretization of $A(\boldsymbol{r}, s, \boldsymbol{v})$ is defined by $F_n(\boldsymbol{r}, s, \boldsymbol{v}) = G_n(R, A(\boldsymbol{r}, s, \boldsymbol{v}))$. The index set of $F_n(\boldsymbol{r}, s, \boldsymbol{v})$ is denoted by $J_n(\boldsymbol{r}, s, \boldsymbol{v})$ and its cardinal number by $j_n(\boldsymbol{r}, s, \boldsymbol{v}) = \operatorname{card}(J_n(\boldsymbol{r}, s, \boldsymbol{v}))$.

Let $r < R \in \mathbb{N}$ and let us define the cut-off function $\theta(r, R, \cdot)$ on $\mathbb{R}$ as a continuous piecewise linear function, such that $\theta(r, R, x) = 1$ for $x \in [-r, r]$, and

Figure 4: The function $\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})$



zero outside of $[-R,R]$. Apparently, $\theta(r,R,\cdot)' = (1/(R-r))\mathbb{1}_{[-R,-r]} - (1/(R-r))\mathbb{1}_{[r,R]}$. Let $\boldsymbol{r}=(r_1,r_2,\ldots,r_d)$ and $\boldsymbol{R}=(R_1,R_2,\ldots,R_d)$. The continuous function $\boldsymbol{x}\mapsto\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v},\boldsymbol{x}) = \prod_{i=1}^d \theta(r_i,R_i,|x_i-v_i|)$ is a cut-off function on $\mathbb{R}^d$ for which $|\partial\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v},\cdot)| \le \max_i(R_i-r_i)^{-1}$.

Discretizations of $\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v},\cdot)$ on $G_n(R)$ are denoted by $\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})$ and they are defined in terms of rougher grids $G_{m-t}\subset G_m\subset G_n$ for which $m-t<m<n$. For each $\boldsymbol{v}\in G_n(R)$, we define three sets

$$\begin{array}{rcl} K(\boldsymbol{r}) &=& S_m(\boldsymbol{p},\boldsymbol{v}) = \{\boldsymbol{x} : \chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v},\boldsymbol{x})=1\},\\ K(\boldsymbol{R}) &=& S_{m-t}(\boldsymbol{p},\boldsymbol{v}) = \mathrm{supp}\,(\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})),\\ K_+(\boldsymbol{R}) &=& K(\boldsymbol{R}) + S_n(\boldsymbol{p},\boldsymbol{0}) \;\;=\;\; \cup_{\boldsymbol{v}\in K(\boldsymbol{R})} S_n(\boldsymbol{p},\boldsymbol{v}), \end{array} \tag{54}$$

which are illustrated in Figure 4. The rectangles (54) have edges $2r_i = p_i 2^{1-m}, 2R_i = p_i 2^{1+t-m}$ and $p_i(2^{1+t-m}+2^{1-n})$ units, respectively. It is important to notice that the grid-functions $\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})$ are defined by sets $S_n(\boldsymbol{p},\boldsymbol{v})$ which are related to the numerical neighbourhoods of constructed schemes. In our next proof we again use the fact that the sets $\mathrm{supp}\,(\max\{a_{ij},0\}), i\neq j$ are strictly separated, so that we can assume that the functions a_{ij} on rectangles (54) do not change sign.

It is easy to verify that $|U_i(p_i)\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})| \le (h(m)\underline{p}(2^t-1))^{-1}$, so that we can write

$$|U_i(p_i)\,\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})| \;\le\; \frac{1}{\underline{p}^2}\,\frac{1}{h(n)^2\,(L-l)^2} \;\le\; \frac{\rho}{|\boldsymbol{R}-\boldsymbol{r}|},$$

where $L=2^{t+n-m}, l=2^{n-m}$, and ρ is a number depending on $\underline{p}=\min p_i(l)$. The sets $F_n(\boldsymbol{r},s,\boldsymbol{v})$ and $F_n(\boldsymbol{R},s,\boldsymbol{v})$ are defined in terms of the sets (54),

$$\begin{array}{rcl} F_n(\boldsymbol{r},s,\boldsymbol{v}) &=& K(\boldsymbol{r})\cap\mathrm{supp}\,(\mathbf{w}_n),\\ F_n(\boldsymbol{R},s,\boldsymbol{v}) &=& K_+(\boldsymbol{R})\cap\mathrm{supp}\,(\mathbf{w}_n). \end{array}$$

The following estimates are used in our next proof:

$$\begin{array}{rcl} \max\limits_{l,i}\; Z(i,\pm p_i(l))\chi(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v}) &\le& \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})},\\ \max\limits_{l,i}\; \big|U_i(p_i(l))\,\big(Z(i,\pm p_i(l))\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})\big)\big| &\le& \dfrac{\rho}{|\boldsymbol{R}-\boldsymbol{r}|}\mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}. \end{array} \tag{55}$$

Now we consider the following auxiliary problem:

$$A_n(D)\,\mathbf{u}_n \;=\; \mathbf{f}_{0n} - \sum_{i=1}^d V_i(p_i)\,\mathbf{f}_{in} \;:=\; \boldsymbol{\mu}_n, \tag{56}$$

where $\mathbf{u}_n$ are grid-solutions and $\mathbf{f}_i$ are grid-functions on $G_n(R,D)$.

LEMMA **6.1** *Let* $\mathbf{u}_n = A_n(D)^{-1}\boldsymbol{\mu}_n$*, where* $\boldsymbol{\mu}_n$ *are defined by (56). There exist positive numbers* c_1, c_2 *and* s_0*, independent of* $n, \boldsymbol{v}$ *such that the inequalities*

$$\begin{aligned}\sum_{i=1}^{d} \big\|\boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})\,U_i(p_i)\mathbf{u}\big\|^2_{R2} \;&\le\; c_1\,\big\|\mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}(\mathbf{u}_n - s)\big\|^2_{R2}\left[1+\frac{1}{|\boldsymbol{R}-\boldsymbol{r}|^2}\right]\\ &+\; c_2\,s^2\sum_{i=0}^{d}\big\|\mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\mathbf{f}_{in}\big\|^2_{Rq}\big(j(\boldsymbol{R},s,\boldsymbol{v})\big)^{1-2/q}\end{aligned} \tag{57}$$

are valid for $s \ge s_0$ *and* $q > 2$.

PROOF: Because of the rule (31) of construction of discretizations for $d > 2$, it is sufficient to prove (57) for two-dimensional problems. Basic schemes are considered first.

The function $w = \max\{u - s, 0\}$ has the discretizations $\mathbf{w}_n$ and $\mathbf{w}_n U_i \mathbf{w}_n = \mathbf{w}_n U_i \mathbf{u}_n$. Now we have to evaluate the forms

$$\begin{aligned} a_n^{(-)}(\chi^2 w, u) \;&=\; \langle\, \boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})^2\mathbf{w}_n \,|\, A_n^{(-)}\,\mathbf{u}_n\,\rangle_R,\\ a_n^{(+)}(\chi^2 w, u) \;&=\; \langle\, \boldsymbol{\chi}_n(\boldsymbol{r},\boldsymbol{R},\boldsymbol{v})^2\mathbf{w}_n \,|\, A_n^{(+)}\,\mathbf{u}_n\,\rangle_R,\end{aligned} \tag{58}$$

in terms of matrices $\mathcal{A}_n^{(\pm)}(i,j,r_i,r_j)$ of Expression (33). It is sufficient to consider the form $a_n^{(-)}$ and one of the terms. For instance, the term involving the matrix $A_n^{(-)}(l)$ is $-\sum_{ij=1}^{2} V_i(p_i)\mathcal{A}_n^{(+)}(i,j,p_i,p_j)\,U_j(p_j)$.

In order to write expressions in this proof in a concise form we omit various indices in the notation. Thus we use $\mathbf{u}, \mathbf{w}, \boldsymbol{\chi}, U_i$ and $\mathcal{A}^{(\pm)}(i,j)$. Apart from this, we need the notations $Z_i\mathbf{f} = Z_n(p_i,i)\mathbf{f}_n$, so that we can write $U_i\mathbf{f}_n = (hp_i)^{-1}(Z_i\mathbf{f} - \mathbf{f})$. In the equality $U_i(\mathbf{g}\mathbf{w}) = (U_i\mathbf{g})Z_i\mathbf{w} + \mathbf{g}U_i\mathbf{w}$ we insert $\mathbf{g} = \boldsymbol{\chi}^2$ to get:

$$U_i\big(\boldsymbol{\chi}^2\mathbf{w}\big) \;=\; \boldsymbol{\chi}^2\,U_i\mathbf{w} + \big(U_i\boldsymbol{\chi}\big)\,\big(Z_i\boldsymbol{\chi}+\boldsymbol{\chi}\big)\,Z_i\mathbf{w}. \tag{59}$$

Also we need

$$\big(Z_i\boldsymbol{\chi}+\boldsymbol{\chi}\big)\,U_j\mathbf{w} \;=\; 2\boldsymbol{\chi}U_j\mathbf{w} + \frac{p_i}{p_j}(U_k\boldsymbol{\chi})Z_j\mathbf{w} - \frac{p_i}{p_j}(U_k\boldsymbol{\chi})\mathbf{w}. \tag{60}$$

In this way the finite differences $U_j\mathbf{w}$ are always multiplied by the grid-function $\boldsymbol{\chi}$ (not with $Z_k\boldsymbol{\chi}$).

We start the proof by an analysis of (58):

$$\begin{aligned}\langle\,\boldsymbol{\chi}^2\,\mathbf{w}\,|\,A_n^{(-)}(l)\,\mathbf{u}\,\rangle_R \;&=\; \sum_{i,j=1}\langle\;\boldsymbol{\chi}\,U_i\mathbf{w}\,|\,\mathcal{A}^{(+)}(i,j)\,\boldsymbol{\chi}\,U_j\,\mathbf{w}\,\rangle_R\\ +\;\sum_{i,j=1}&\langle\,(Z_i\mathbf{w})\,U_i\boldsymbol{\chi}\,|\,\mathcal{A}^{(+)}(i,j)\,\big(Z_i\boldsymbol{\chi}+\boldsymbol{\chi}\big)\,U_j\,\mathbf{w}\,\rangle_R.\end{aligned}$$

The first term on the right hand side can be estimated from bellow as a consequence of the strict ellipticity (45) of Proposition 5.2. The left hand side of this equality is equal to $\langle\,\boldsymbol{\chi}^2\mathbf{w}, |\boldsymbol{\mu}\,\rangle_R$, so that we can write:

$$\begin{aligned}\underline{M}\sum_{i=1}\big\|\boldsymbol{\chi}\,U_i\mathbf{w}\big\|^2_{R2} \;&\le\; \langle\,\boldsymbol{\chi}^2\,\mathbf{w}\,|\,\boldsymbol{\mu}\,\rangle_R\\ -\;\sum_{i,j=1}&\langle\,Z_i\mathbf{w}\,U_i\boldsymbol{\chi}\,|\,\mathcal{A}^{(+)}(i,j)\,\big(Z_i\boldsymbol{\chi}+\boldsymbol{\chi}\big)\,U_j\,\mathbf{w}\,\rangle_R.\end{aligned} \tag{61}$$

The two terms on the right hand side are denoted by $T_i, i = 1, 2$, respectively. Let us estimate them from above. The representation of $\boldsymbol{\mu}$, in (56), is inserted into the first term to get:

$$|T_1| \;=\; |\langle\,\boldsymbol{\chi}^2\,\mathbf{w}\,|\,\boldsymbol{\mu}\,\rangle_R| \;\le\; |\langle\,\boldsymbol{\chi}^2\,\mathbf{w}|\mathbf{f}_0\,\rangle_R| + \sum_{i=1}^{2}|\langle\,U_i(\boldsymbol{\chi}^2\,\mathbf{w})|\mathbf{f}_i\,\rangle_R|.$$

First (59) and then (60) are utilized in the last two terms on right hand side. The functions $\boldsymbol{\chi}, Z_k\boldsymbol{\chi}$ are estimated from above by (55) and then the CSB inequality and the inequality $ab \leq (\varepsilon/2)a^2 + (1/2\varepsilon)b^2$ with a convenient choice of ε, is applied to all the terms on the right hand side.

$$\begin{aligned} |T_1| \;&\leq\; \varepsilon \sum_{i=1}^{2} \left\| \boldsymbol{\chi}\, U_i \mathbf{w} \right\|_{R2}^2 + b_1\Big(1 + \frac{1}{|\boldsymbol{R}-\boldsymbol{r}|^2}\Big) \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{w} \right\|_{R2}^2 \\ &+\; \Big(1 + \frac{1}{4\varepsilon}\Big) \sum_{i=0}^{2} \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{f}_i \right\|_{R2}^2, \end{aligned}$$

where b_1 does not depend on n, ε. It is important to notice that the first term on the right hand side of this inequality and the left hand side of (61) involve the same norms, and can therefore be subtracted. Now we use the Hölder inequality $\left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})} \mathbf{f}_i \right\|_{R2}^2 \leq \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})} \mathbf{f}_i \right\|_{Rq}^2 j(\boldsymbol{R}, s, \boldsymbol{v})^{1-2/q}$ and get the first intermediate result:

$$\begin{aligned} |T_1| \;&\leq\; \varepsilon \sum_{i=1}^{2} \left\| \boldsymbol{\chi}\, U_i \mathbf{w} \right\|_{R2}^2 + b_1\,\Big(1 + \frac{1}{|\boldsymbol{R}-\boldsymbol{r}|^2}\Big) \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{w} \right\|_{R2}^2 \\ &+\; b_2 \sum_{i=0}^{2} \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{f}_i \right\|_{Rq}^2 j(\boldsymbol{R}, s, \boldsymbol{v})^{1-2/q}, \end{aligned}$$

where b_2 depends on $1/\varepsilon$.

The quantity T_2 can be estimated analogously. Let us use (60) in Expression (61) and apply the same technique as in the previous case. We get:

$$|T_2| \;\leq\; \varepsilon \sum_{i=1}^{2} \left\| \boldsymbol{\chi}\, U_i \mathbf{w} \right\|_{R2}^2 + b_3\, \frac{1}{|\boldsymbol{R}-\boldsymbol{r}|^2} \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{w} \right\|_{R2}^2,$$

where b_3 depends on $\overline{M}, 1/\varepsilon$. Thus the inequality (61) implies the following result:

$$\begin{aligned} \big(\underline{M} - 2\varepsilon\big) \sum_{i=1}^{2} \left\| \boldsymbol{\chi}\, U_i \mathbf{u} \right\|_{R2}^2 \;&\leq\; b_4\,\Big(1 + \frac{1}{|\boldsymbol{R}-\boldsymbol{r}|^2}\Big) \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{w} \right\|_{R2}^2 \\ &+\; b_2 \sum_{i=0}^{2} \left\| \mathbb{1}_{F(\boldsymbol{R},s,\boldsymbol{v})}\, \mathbf{f}_i \right\|_{Rq}^2 j(\boldsymbol{R}, s, \boldsymbol{v})^{1-2/q}, \end{aligned}$$

which is equivalent to the assertion of lemma.

Now we consider the extended schemes. Let the discretizations $A_n(D)$ of (57) be constructed by extended schemes and let $\mathbf{u}_n$ be the corresponding solution. Then we have $a_n(v, u) = \langle\, \mathbf{v}_n | \boldsymbol{\mu}_n \,\rangle$, where $a_n(v, u)$ is defined in (34). Let us define $\mathbf{v}_n = \boldsymbol{\chi}^2 \mathbf{w}$ and let its image $v(n) = \Phi_n \mathbf{v}_n$ be inserted into the expression $a_n(v, u)$. If we apply (59) to the function $v(n)$ we get:

$$a_n(v(n), u) \;=\; a_n(\chi; w, w) + \sum_s b_n\big(s, \boldsymbol{\chi}, \mathbf{w}\big), \tag{62}$$

where the terms on the right hand side are defined as follows. The form $a_n(\chi; w, w)$ is the sum of forms (47) and (48) in which the factors $a_{ij}(\boldsymbol{x} + \boldsymbol{t}^{(\pm)}(\boldsymbol{r}))$ are replaced with the factors $\boldsymbol{\chi}(\boldsymbol{x})^2 a_{ij}(\boldsymbol{x} + \boldsymbol{t}^{(\pm)}(\boldsymbol{r}))$. Each $b_n(s, \boldsymbol{\chi}, \mathbf{w})$ is a form which is a first degree polynomial in $U_i(p_i(l))\mathbf{w}_n$, with various $l, i, p_i(l)$, resulting from the application of the rule (59). As in the corresponding proof of Proposition 5.2, the quantity $a_n(\chi; w, w)$ can be estimated from bellow,

$$a_n(\chi; w, w) \;\geq\; \omega(a) \sum_{i=1}^{2} \left\| \boldsymbol{\chi}\, U_i \mathbf{w} \right\|_2^2.$$

An attempt to estimate the remaining terms on the right hand side in (62) from above as in the first part of proof would lead to an unresolvable problem. Among the resulting terms there would be $\varepsilon \sum_i \bracevert \boldsymbol{\chi}_n U_i \mathbf{w}_n \bracevert_2^2$ as well as $\varepsilon \sum_i \bracevert \boldsymbol{\chi}_n U_i(p_i) \mathbf{w}_n \bracevert_2^2$. The former one can be moved to the left hand side of (62) and subtracted from $\omega(a) \sum_i \bracevert \boldsymbol{\chi}_n U_i \mathbf{w}_n \bracevert_2^2$. Unfortunately, this cannot be done with the latter one. Therefore, before finding upper bounds on $b_n(s, \boldsymbol{\chi}, \mathbf{w})$ we have to apply another version of (60):

$$\begin{aligned}\big(\boldsymbol{\chi}_n + Z_n(p_i, i)\boldsymbol{\chi}_n\big) U_j(p_j)\mathbf{w}_n &= 2\tilde{\boldsymbol{\chi}}_n U_j(p_j)\mathbf{w}_n + \\ \frac{p_i}{p_j}\big(U_i(p_i)\tilde{\boldsymbol{\chi}}_n\big)\big(Z_n(p_j, j) - I\big)\mathbf{w}_n &+ \frac{1}{p_j}\Big[\big(I + Z_n(p_i, i)\big)\big(\boldsymbol{\chi}_n - \tilde{\boldsymbol{\chi}}_n\big)\Big]\big(Z_n(p_j, j) - I\big)\mathbf{w}_n.\end{aligned}$$

If these expressions are applied, then estimates from above of $b_n(s, \boldsymbol{\chi}, \mathbf{w})$ contain the term $\varepsilon \sum_i \bracevert \tilde{\boldsymbol{\chi}}_n U_i(p_i) \mathbf{w}_n \bracevert_2^2$. Let us choose $\tilde{\chi}(\boldsymbol{x}) = \chi(\boldsymbol{r}, \boldsymbol{R}, \boldsymbol{v}, \boldsymbol{x})\chi(\boldsymbol{r}, \boldsymbol{R}, \boldsymbol{v}, \boldsymbol{x} + h(p_1\boldsymbol{e}_1 +_2 \boldsymbol{e}_2))$. Then

$$\operatorname{supp}(\tilde{\chi}) + C_n(\boldsymbol{p}, \mathbf{0}) \;=\; K(\boldsymbol{R}),$$

and $(\tilde{\boldsymbol{\chi}}_n U_j(p_j)\mathbf{w}_n)_{\boldsymbol{k}}$ is a linear combination of $(U_j(1)\mathbf{w}_n)_{\boldsymbol{l}}$ such that $h\boldsymbol{l} \in K(R)$. Now we have

$$\varepsilon \sum_i \bracevert \tilde{\boldsymbol{\chi}}_n U_i(p_i) \mathbf{w}_n \bracevert_2^2 \;\le\; \varepsilon \sum_i \bracevert \boldsymbol{\chi}_n U_i(1) \mathbf{w}_n \bracevert_2^2$$

and the difficulty regarding $\bracevert \boldsymbol{\chi}_n U_i(p_i) \mathbf{w}_n \bracevert_2^2$ with various $p_i \neq 1$ is removed. The grid-functions $h_{-1}(\boldsymbol{\chi}_n - \tilde{\boldsymbol{\chi}}_n)$ have estimates from above as in (55). In this way we get (57) again. **QED**

The function $w(n) = \max\{u(n) - s, 0\}$ and the grid-function $\mathbf{w}_n$ which is componentwisely by $w_{\boldsymbol{k}} = \max\{u_{\boldsymbol{k}} - s, 0\}$, are related by the expression $w(n) = \Phi_n(R)\mathbf{w}_n$. The index set of grid-knots of $G_n(R)$ in the (closed) set $S_m(\boldsymbol{p}, \boldsymbol{v})$ is denoted by $I(m, \boldsymbol{p}, \boldsymbol{v})$. The restriction of $w(n)$ to the set $S_m(\boldsymbol{p}, \boldsymbol{v})$ and the function $w(n, I) = \sum_{\boldsymbol{k} \in I(m, \boldsymbol{p}, \boldsymbol{v})} w_{\boldsymbol{k}} \psi_{\boldsymbol{k}}$ are not equal. They are equal on the set $S_m(\boldsymbol{p}, \boldsymbol{v})$ and the following rough estimate is valid elsewhere:

$$\begin{aligned}\|w(n, I)\|_2^2 &= \|\mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})}\, w(n, I)\|_2^2 \;+\; \|(1 - \mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})})\, w(n, I)\|_2^2 \\ &\le 2^d\, \|\mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})}\, w(n, I)\|_2^2 \;=\; 2^d\, \|\mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})}\, w(n)\|_2^2.\end{aligned}$$

Now we combine the result of Theorem 3.1, *i.e.*:

$$(1 - \sigma^2) \bracevert \mathbb{1}_{I(m, \boldsymbol{p}, \boldsymbol{v})}\, \mathbf{w}_n \bracevert_{R2}^2 \;\le\; \|w(n, I)\|_2^2 \;\le\; \bracevert \mathbb{1}_{I(m, \boldsymbol{p}, \boldsymbol{v})}\, \mathbf{w}_n \bracevert_{R2}^2$$

and the previous inequality in order to get:

$$\bracevert \mathbb{1}_{I(m, \boldsymbol{p}, \boldsymbol{v})}\, \mathbf{w}_n \bracevert_{R2}^2 \;\le\; \frac{1}{1 - \sigma^2}\, \|w(n, I)\|_2^2 \;\le\; \frac{2^d}{1 - \sigma^2}\, \|\mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})}\, w(n)\|_2^2 \qquad (63)$$

It remains to compare the functions $\partial_i w(n)$ and $U_i(p_i)\mathbf{w}_n$. By Expression (26) we easily prove:

$$\begin{aligned}\|\mathbb{1}_{S_m(\boldsymbol{p}, \boldsymbol{v})}\, \partial_i w(n)\|_2^2 \;\le\; \|\textstyle\sum_{\boldsymbol{k} \in I(m, \boldsymbol{p}, \boldsymbol{v})}\, \partial_i w(n)\|_2^2 \;&\le\; h^d \bracevert \mathbb{1}_{I(m, \boldsymbol{p}, \boldsymbol{v})}\, \big(U_i(p_i)\mathbf{w}_n\big)_{\boldsymbol{k}} \bracevert_{R2}^2 \\ &\le\; h^d \bracevert \boldsymbol{\chi}_n(\boldsymbol{r}, \boldsymbol{R}, \boldsymbol{v})\, U_i(p_i)\mathbf{w}_n \bracevert_{R2}^2.\end{aligned} \qquad (64)$$

Let us multiply (57) by h^d, use (64) and afterwards (63). The result is:

$$\begin{aligned}\textstyle\sum_{i=1}^d \|\mathbb{1}_{A(\boldsymbol{r}, s, \boldsymbol{v})}\, \partial_i w(n)\|_2^2 \;&\le\; \frac{2^d\, c_1}{1 - \sigma^2}\, \|\mathbb{1}_{A(\boldsymbol{R}, s, \boldsymbol{v})}\, w(n)\|_2^2 \left(1 + \frac{1}{|\boldsymbol{R} - \boldsymbol{r}|}\right) \\ &+\; c_2\, s^2 \left(h^{2d/q} \sum_{i=0}^d \bracevert \mathbf{f}_{in} \bracevert_{Rq}^2\right) a(\boldsymbol{R}, s, \boldsymbol{v})^{1 - 2/q}.\end{aligned} \qquad (65)$$

In this way we obtain an important property of grid-solutions to Problems (56).

COROLLARY **6.1** *Let $D \subset \mathbb{R}^d$ be a bounded domain with a Lipshitz boundary and $q > d$. If $h^d \sum_{i=0}^d \bracevert \mathbf{f}_{in} \bracevert_{Rq}^q$ are bounded by a number uniformly with respect to n, then the grid-solutions $\mathbf{u}_n$ on $G_n(R, D)$ to (56) are also bounded uniformly with respect to n.*

PROOF: One has to compare (53) and inequality (65). **QED**

6.2 Weak consistency and convergence

As in the case of W_2^1-approach, the convergence in L_p-spaces is proved by utilizing certain kind of consistency. This weaker consistency is defined in terms of sequences similar to (40):

$$\begin{array}{lclcl} \mathfrak{V}(p,c) & = & \{v(n) \,:\, n \in \mathbb{N},\ h^d \sum_i \|U_i \mathbf{v}_n\|_p^p \leq c\} & \subset & \cup_n E_n(\mathbb{R}^d), \\ \mathfrak{U}_0 & = & \{\hat{u}(n) \,:\, n \in \mathbb{N},\ u \in C_0^{(1)}(\mathbb{R}^d)\} & \subset & \cup_n E_n(\mathbb{R}^d). \end{array} \tag{66}$$

DEFINITION 6.1 (Weak consistency) *We say that forms* $a_n(\cdot,\cdot)$ *on* $E_n(\mathbb{R}^d) \times E_n(\mathbb{R}^d)$ *are weakly consistent with the form (3) if*

$$a(v,u) \;=\; \lim_n h^d\, a_n(v(n), \hat{u}(n))$$

is valid for any weakly convergent $\mathfrak{V}(p,c), p \in [1,\infty), c > 0$ *and any* $\mathfrak{U}_0$ *of (66).*

LEMMA **6.2** *Let a sequence of matrices* $\{A_n : n \in \mathbb{N}\}$ *be constructed by basic or extended schemes, and let* $\{a_n(\cdot,\cdot) : n \in \mathbb{N}\}$ *be the corresponding sequence of discretizations of (3). Then the forms* a_n *are weakly consistent with the form (3).*

Proof: The present proof and proof of Lemma 5.1 are the same up to the equality (44). The obtained equality must be estimated now by the Hölder inequality:

$$\Big|\gamma_n(l, v(n), u(n)) \;-\; \theta_n(l, v(n), u(n))\Big| \;\leq$$
$$\overline{p}\, h^{d/p} \|U_i(r_i)\mathbf{v}_n\|_{Rp}\, h^{d/q} \max\{\|\big(Z_n(r_j,j) - I\big)U_j(r_j)\hat{\mathbf{u}}_n\|_{Rq} \,:\, j = 1,2,\ldots,d\}.$$

Thus we have to consider $\|\big(Z_n(r_j,j) - I\big)U_j(r_j)\hat{\mathbf{u}}_n\|_{Rq}$ for a fixed $j \in \mathbb{N}$ and prove that this quantity converges to zero as $n \to \infty$. It is easy to get the following expression:

$$\Big(\big(Z_n(r_j,j) - I\big)\, U_j(r_j)\hat{\mathbf{u}}_n\Big)_{\boldsymbol{k}} \;=\; \frac{1}{\|\psi_{\boldsymbol{k}}\|_1} \big(\psi_{\boldsymbol{k}} \,|\, (Z(th\boldsymbol{e}_j) - I\big) \exists_j(r_j h) u\big) \;=\; \big(\hat{\mathbf{w}}_n(r_j)\big)_{\boldsymbol{k}},$$

where the function $w(r_j) = \big(Z(th\boldsymbol{e}_j) - I\big) \exists_j(r_j h) u$ is continuous and converges pointwisely to zero as $h \to 0$. By (ii) of Lemma 3.6 we get

$$h^{d/q}\, \|\hat{\mathbf{w}}_n(r_j)\|_{Rq} \;\leq\; \|w(r_j)\|_q \;\; \underset{h \to 0}{\longrightarrow} \;\; 0.$$

QED

THEOREM **6.1** *Let* $D \subset \mathbb{R}^d$ *be a bounded domain with a Lipshitz boundary, and let* $\mathfrak{U} = \{u(n) : n \in \mathbb{N}\} \subset E_n(R,D)$ *be a sequence of approximate solutions,* $u(n) = \Phi_n(R)\mathbf{u}_n$*, where* $\mathbf{u}_n$ *are grid-solutions to (52). For each* $p \in [1, d/(d-1))$ *there exists a subset* $J(p) \subset \mathbb{N}$ *such that* $\mathfrak{U} = \{u(n) : n \in J(p)\}$ *converges to the unique solution to (5) strongly in* $L_p(D)$ *and weakly in* $\dot{W}_p^1(D)$*.*

Proof: The proof is split into two steps. In the first step we have to prove that for each $p \in (1, d/(d-1))$ there exits $u \in \dot{W}_p^1(D)$ and a subset $J(p) \subset \mathbb{N}$ such that

$$\begin{array}{lclcl} u & = & w\text{–}\lim\limits_{n \in J(p)} u(n) & \in & \dot{W}_p^1(D), \\ u & = & s\text{–}\lim\limits_{n \in J(p)} u(n) & \in & L_p(D). \end{array}$$

In the second step of the proof we demonstrate that u coincides with the solution u^* of the original problem (5).

Because of $D \subset S_1(\boldsymbol{t}, \boldsymbol{v})$, for some $\boldsymbol{t}, \boldsymbol{v}$ we have $\|u\|_p \leq (\max_k t_k)\,(\max_k \|\partial_k u\|_p)$ so that we can consider only $\partial_1 u(n)$ and the corresponding $U_1(p_1)\mathbf{u}_n$. By (26) and estimating the function $\psi_{\boldsymbol{k}'}$ by the indicator of its support we get:

$$|\partial_1 u(n)| \;\leq\; 2^d \sum_{\boldsymbol{k} \in I_n(R)} \Big|\big(U_1(p_1)\mathbf{u}_n\big)_{\boldsymbol{k}}\Big|\, \mathbb{1}_{C_n(\boldsymbol{p},\boldsymbol{v})},$$

Figure 5: Domains for Examples 7.1 and 7.2



Example 7.1 Example 7.2

where $\boldsymbol{v} = h\boldsymbol{k}$. Hence

$$\|\partial_1 u(n)\|_p^p \;\leq\; 2^{dp}\, h^d \,\boldsymbol{|} U_1(p_1)\mathbf{u}_n \boldsymbol{|}_{Rp}^p .$$

It remains to demonstrate that the right hand side is bounded uniformly with respect to n. We can write

$$\boldsymbol{|} U_1(p_1)\mathbf{u}_n \boldsymbol{|}_{Rp} \;=\; \langle\, \mathbf{v}_n \,|\, U_1(p_1)\, \mathbf{u}_n \,\rangle_R ,$$

where $v_{\boldsymbol{k}} = h^{-d/q}\mathrm{sign}(U_1(p_1)(\mathbf{u}_n)_{\boldsymbol{k}})|U_1(p_1)(\mathbf{u}_n)_{\boldsymbol{k}}|^{p-1}/\boldsymbol{|} U_1(p_1)\mathbf{u}_n \boldsymbol{|}_{Rp}^{p-1}$. Then $h^{d/q}\boldsymbol{|}\mathbf{v}_n\boldsymbol{|}_{Rq} = 1$. Therefore,

$$\boldsymbol{|} U_1(p_1)\mathbf{u}_n \boldsymbol{|}_{Rp} \;=\; -\,\langle\, V_1(p_1)\mathbf{v}_n \,|\, \mathbf{u}_n \,\rangle_R \;=\; -\,\langle\, A_n^{-1}\left(V_1(p_1)\mathbf{v}_n\right) |\, \boldsymbol{\mu}_n \,\rangle_R .$$

By Corollary 6.1 we get $\boldsymbol{|} A_n^{-1} V_1(p_1)\mathbf{v}_n \boldsymbol{|}_{R\infty} \leq \beta$ with some β depending on D and the bounds of $h^d\boldsymbol{|}\mathbf{v}_n\boldsymbol{|}_{Rq}^q$. The final result is $h^{d/p}\boldsymbol{|} U_i(p_i)\mathbf{u}_n \boldsymbol{|}_{Rp} \leq \beta$.

The unique solution to (5) is denoted by u^*. For $p \in (1, d/(d-1))$, $v \in \dot{C}^{(1)}(\overline{D})$ and the corresponding $\hat{\mathbf{v}}_n, \hat{v}(n) = \Phi_n(R)\hat{\mathbf{v}}_n$, the following relations are valid:

$$\begin{array}{ll}
1) & v \;=\; s\text{–}\lim_n \hat{v}(n) \;\in\; \dot{W}_q^1(D), \qquad 1/p + 1/q = 1, \\
2) & h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, \boldsymbol{\mu}_n \,\rangle_R \;=\; \langle\, \hat{v}(n) \,|\, \mu \,\rangle \;\rightarrow\; \langle\, v \,|\, \mu \,\rangle, \\
3) & \lim_n a(\hat{v}(n), u(n)) \;-\; \lim_n h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, A_n(D)\mathbf{u}_n \,\rangle_R \;=\; 0.
\end{array}$$

The inclusion 1) follows from Lemma 3.4 and the uniform continuity of ∂v_1 on $\overline{D}$. Although this inclusion is plausible, some technical details are needed in order to transfer ∂_1 from $\partial_1\hat{v}(n)$ to $(\widehat{\partial_1 v})(n)$. The relation 2) follows from $v \in \dot{C}(\overline{D})$ and the construction of $\boldsymbol{\mu}_n$. The identity 3) is the weak consistency which is proved in Lemma 6.2. Notice that the pair $\hat{v}, u$ in this proof and the pair $v, \hat{u}$ of Lemma 6.2 have roles interchanged. Now we have

$$\begin{array}{rcl}
a(v, u) & = & a(v, u) \;-\; a(\hat{v}(n), u(n)) \\
& + & a(\hat{v}(n), u(n)) \;-\; h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, A_n(D)\, \mathbf{u}_n \,\rangle_R \\
& + & h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, A_n(D)\, \mathbf{u}_n \,\rangle_R \;-\; h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, \boldsymbol{\mu}_n \,\rangle_R \\
& + & h^d\, \langle\, \hat{\mathbf{v}}_n \,|\, \boldsymbol{\mu}_n \,\rangle_R .
\end{array}$$

The right hand sides in the first three rows are either zero or converge to zero as $n \to \infty$. Thus $a(v, u) = \langle\, v|\mu\,\rangle$. Because $\dot{C}^{(1)}(\overline{D})$ is dense in $\dot{W}_p^1(D), p \in (1, d/(d-1))$ we get $u = u^*$. However $u - u^* = 0 \in \dot{W}_p^1(D)$ implies $u - u^* = 0 \in \dot{W}_1^1(D)$. **QED**

7 Numerical examples in two dimensional case

We test the numerical methods on examples for which we know solutions in the closed form. In the first example we solve the problem (4) with $a_{12} = 0$ and a measure on the right hand side, and in the second exmple we have a piecewise constant a_{12} and a non-trivial application of the extended scheme.

EXAMPLE **7.1** Let the matrix-valued function $\boldsymbol{x} \mapsto a(\boldsymbol{x})$ be defined on $\mathbb{R}^2$ by

$$a(\boldsymbol{x}) \;=\; 10^{-4} \times \left\{ \begin{array}{ll} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} & \text{for} \quad x_1 < 0.5, \\ \begin{bmatrix} \sigma^2 & 0 \\ 0 & 1 \end{bmatrix} & \text{for} \quad x_1 \geq 0.5, \end{array} \right.$$

and let us consider the elliptic differential operator $A(\boldsymbol{x}) = -\sum_{ij=1}^{2} \partial_i a_{ij}(\boldsymbol{x})\partial_j$ on $\mathbb{R}^2$. For $\boldsymbol{t} = (0.5, 0.5)$ the differential equation $A(\boldsymbol{x})F(\boldsymbol{x}) = \delta(\boldsymbol{x} - \boldsymbol{t})$ on $\mathbb{R}^2$ has a solution

$$F(\boldsymbol{x}) \;=\; -c \left\{ \begin{array}{ll} \ln\left((x_1 - 0.5)^2 + (x_2 - 0.5)^2\right) & \text{for} \quad x_1 \;<\; 0.5, \\ \left[\ln\left((x_1 - 0.5)^2) + \sigma^2 (x_2 - 0.5)^2\right) - \ln(\sigma^2)\right] & \text{for} \quad x_1 \;\geq\; 0.5, \end{array} \right.$$

where

$$c \;=\; \Big(2\pi\left(1 + \sigma\right)\Big)^{-1}.$$

The function $F(\cdot)$ is continuous on $\mathbb{R}^2 \setminus \{\boldsymbol{t}\}$ and its first partial derivatives have a jump at $x_1 = 1/2$. Let $D = (0, 1)^2 \subset \mathbb{R}^2$ as illustrated in Figure 5. In this example we consider the boundary value problem $A(\boldsymbol{x})u(\boldsymbol{x}) = \delta(\boldsymbol{x} - \boldsymbol{t})$ with the nonhomogeneous boundary conditions $u|\partial D = F|\partial D$. Obviously, the solution u coincides with F on D.

In numerical calculations we used $\sigma^2 = 10$. Hence, we have two sets for the discretization procedure 4.1, $D_1 = \{\boldsymbol{x} \in \mathbb{R}^2 : x_1 < 0\}$ and $D_2 = \{\boldsymbol{x} \in \mathbb{R}^2 : x_2 \geq 0\}$. The extended scheme was utilized for the discretization of $A(\boldsymbol{x})$. Discretization of $\mathbb{R}^2$ is realized by grid-knots $\boldsymbol{x} = hk_1\boldsymbol{e}_1 + hk_2\boldsymbol{e}_2$, $h = 1/400$, so that D is discretized by 399×399 grid-knots. The system matrix A_n of (39) has the order 399 and the linear system is solved iteratively: if $K_n = \mathrm{diag}(A_n)$ and $Q_n = \mathrm{diag}(A_n) - A_n$, then $Q_n \geq 0$ and

$$A_n^{-1} \;=\; \sum_{k=0}^{\infty} \left(K_n^{-1} Q_n\right)^k K_n^{-1}. \tag{67}$$

Let $\mathbf{u}_n(r)$ be the approximation of $\mathbf{u}_n$ after r iterations. By taking the stopping criteria to be $\|\mathbf{u}_n(r{+}1) - \mathbf{u}_n(r)\|_1 < 10^{-9}$, we have found that the iteration terminates after $r = 182000$ iterations. Then we compared the numerical approximations $\mathbf{u}_n(210)$ and values of the solution in $l_\infty(G_n(D) \setminus \{\boldsymbol{t}\})$ and $l_1(G_n(D) \setminus \{\boldsymbol{t}\})$–norms, where the set $G_n(D) \setminus \{\boldsymbol{t}\}$ is chosen naturally instead of the set G_n because the solution u of this example is not defined at $\boldsymbol{t}$. Let the corresponding norms be denoted by $\|\cdot\|_p'$, $p = 1, \infty$. The solutions are compared according to the relative error:

$$\varepsilon_{p,rel} \;=\; \frac{\|\mathbf{u}^* \;-\; \mathbf{u}_n(210)\|_p'}{\|\mathbf{u}^*\|_p'}, \qquad p = 1, \infty, \tag{68}$$

where $\left(\mathbf{u}^*\right)_{\boldsymbol{k}} = u(h\boldsymbol{k})$. We obtained $\varepsilon_{1,rel} = 0.003371$ and $\varepsilon_{\infty,rel} = 1.6254$. The value $\varepsilon_{\infty,rel} = 1.6254$ is realized at the grid-knot $\boldsymbol{t} - h\boldsymbol{e}_1$, i.e. at one of the nearest grid-knots to the singular point $\boldsymbol{t}$ of solution. Actually, the difference between two solutions at this grid-knot is 0.0075. Let us mention that the respective error $\varepsilon_{1,rel} = 0.267$ is obtained with the grid-step $h = 1/200$.

EXAMPLE **7.2** Next, let us consider a differential operator $A = -\sum_{ij=1}^{2} \partial_i a_{ij} \partial_j$ with the diffusion tensor

$$a \;=\; \begin{bmatrix} \sigma^2 & \alpha(\boldsymbol{x}) \\ \alpha(\boldsymbol{x}) & 1 \end{bmatrix}, \quad \alpha(\boldsymbol{x}) \;=\; \rho \mathbb{1}_{D_0}(\boldsymbol{x}), \quad \rho^2 < \sigma^2,$$

where σ^2 is a positive number, ρ is a real number and $D_0 = (1/4, 3/4)^2$ (see Figure 5). The function $\boldsymbol{x} \mapsto u^*(x_1, x_2) = x_1 x_2$ is the unique solution to the boundary value problem

$$\begin{aligned} \big(Au\big)(\boldsymbol{x}) \;&=\; \mu(\boldsymbol{x}) \quad \text{for} \quad \boldsymbol{x} \in D, \\ u|\partial D \;&=\; u^*|\partial D, \end{aligned}$$

where

$$\mu(\boldsymbol{x}) \;=\; 2\,\rho\,\mathbb{1}_{D_0}(\boldsymbol{x}) \,+\, \frac{\rho}{4}\Big[\delta\Big(x_1-\frac{1}{4}\Big)-3\delta\Big(x_1-\frac{3}{4}\Big)+\delta\Big(x_2-\frac{1}{4}\Big)-3\delta\Big(x_2-\frac{3}{4}\Big)\Big].$$

The set $\mathbb{R}^2$ is discretized as in the previous example. The sets $G_n(2) = \overline{D_0} \cap G_n$ and $G_n(1) = G_n \setminus G_n(0)$ define a partition of G_n. Let $G_n(1,D) = G_n(1) \cap D$. Then $G_n(2), G_n(1,D)$ is a partition of $G_n(D)$ to be used in constructions of numerical grids and approximate solutions.

To demonstrate the efficiency of the extended schemes we choose $\sigma^2 = 10, \rho = 2$ and the scheme parameters $r_1 = 1, r_2 = 3$ as illustrated in Figure 3. These values of parameters ensure the compartmental structure of the system matrix. The linear system (52) is solved iteratively as in the previous example. Let $\mathbf{u}_n(r)$ be the approximation of $\mathbf{u}_n$ after r iterations. By taking the stopping criteria to be $\|\mathbf{u}_n(r+1)-\mathbf{u}_n(r)\|_1 < 10^{-9}$, we have found that the iteration terminates after $r = 210$ iterations. Then we compared the numerical approximations $\mathbf{u}_n(210)$ and values of the solution in $l_\infty(G_n(D))$ and $l_1(G_n(D))$–norms. We obtained $\varepsilon_{1,rel} = 0.8 \times 10^{-5}$. In addition, $\|\mathbf{u}_n^* - \mathbf{u}_n(210)\|_{\infty,rel} = 0.4 \times 10^{-2}$ is realized at the grid-knot with coordinates $\boldsymbol{x} = (0.75, 0.75)$.

8 DISCUSSION AND CONCLUSION

Attempts to discretize a second order elliptic differential operator by matrices with compartmental structure go back to the work of Motzkyn and Wasov [MW]. Their construction is based on rotations, which can be avoided in 2-dimensional cases by using extended schemes which we propose in this work. These can be used to get monotone schemes for any 2-dimensional problem with the second order elliptic operator, in divergence or non-divergence form. For operators in divergence form one has to use the construction of system matrix described in Section 4, while for operators in non-divergence form system matrices are of a simpler structure. Entries of system matrices are linear functions of $a_{ij}(\boldsymbol{v})$, where $\boldsymbol{v}$ is the considered grid-knot.

In the case of dimension $d \geq 3$ the construction of Section 4 does not always produce monotone schemes. An additional condition on the matrix-valued function $\boldsymbol{x} \mapsto \{a_{ij}(\boldsymbol{x})\}_{11}^{dd}$ which is described in Assumption 4.1 assures the compartmental structure of matrices A_n.

LEMMA **8.1** *Let A_n on G_n be consistent discretizations of the differential operator $A(\boldsymbol{x}) = -\sum_{ij}\partial_i a_{ij}(\boldsymbol{x})\partial_j$ and let the entries $\big(A_n\big)_{\boldsymbol{kl}}$ be linear combinations of $a_{ij}(\boldsymbol{z})$ with some $\boldsymbol{z} \in S_n(\boldsymbol{p}(l), h\boldsymbol{x})$. Then A_n can possess the compartmental structure iff the tensor-valued function $\hat{a}$ is strictly positive definite on $\mathbb{R}^d$.*

PROOF: For a constant diffusion tensor we prove easily this lemma by demanding the following property at each grid-knot:

$$\big(A_n \mathbf{f}_n\big)_{\boldsymbol{k}} \;=\; A(\boldsymbol{x}) f(\boldsymbol{x}), \quad \boldsymbol{x} = h\boldsymbol{k},$$

where $\boldsymbol{x} \mapsto f(\boldsymbol{x})$ is any second degree polynomial in variables x_i. This property is equivalent to the consistency. To prove the necessary part, we assume that $\hat{a}(\boldsymbol{x}_0)$ is negative definite and A_n have the compartmental structure. Then the same assertion must hold for a tensor-valued function which is constant locally at $\boldsymbol{x}_0$. This contradicts the first step. **QED**

Hence, if the matrix-valued function $\hat{a}$ is indefinite the only way out is to use rotations. Unfortunately undesired features may appear; in a subdomain $D_l \subset D$ where a rotation is required, some larger diagonal submatrices can be reducible. This causes the surfaces $u(n,\boldsymbol{x}) = \mathit{const}$ to have a saw-like behavior. In this case an averaging procedure of grid-solutions helps to obtain reasonable results.

A different approach to the construction of monotone schemes for a class of elliptic operators was presented in Samarskii *et al.* [SMMM]. They constructed schemes with various grid-steps along coordinate axis which we called basic schemes in our analysis. In this way they obtained monotone schemes for a wide class of elliptic operators.

References

[Ba] R. F. Bass, *Diffusions and Elliptic operators*, Springer, N. Y. 1977

[BPV] I. Babuska, M. Prager and E. Vitasek *Numerical Processes in Differential Equations*, SNTL-Publishers, Prague, 1966

[Cl] S. Clain, *Finite element approximations for the Laplace equation with a right-hand side measure*, Math. Models Methods Appl. Sci, **5**, 713–719, 1996

[BO] G. Bottaro, P. Oppezzi, *Elliptic equations in divergence form, with right hand side a measure, on unbounded domains*, Ann. Mat. Pura Appl., 163, 223–246, 1993

[BS] G. Barles and P. E. Sougandis, *Convergence of approximation schemes for fully nonlinear second order equations*, Asymptotic Analysis Vol 5., 271-283, 1991

[Gi] M. Giaquinta, *Introduction to Regularity Theory for Nonlinear Elliptic Systems*, Birkhäuser, Basel, Lectures in Mathematics, 1993

[GT] D. Gilberg and N. S. Trudinger, *Elliptic Partial Differential Equations*, Springer, Berlin, 1983

[LU] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Partial Differential Equations*, Academic Press, N.Y., 1968

[LR1] N. Limić and M. Rogina, *Explicit stable methods for second order parabolic systems*, Math. Commun. Vol 5, 97–115, 2000

[LR2] N. Limić and M. Rogina, *Numerical solutions of Dirichlet problem for elliptic operator in divergence form with a right-hand side measure*, J. Comput. Appl. Math., 493–516, 2004

[MW] T. S. Motzkyn and W. Wasov, *On the approximation of linear elliptic differential equations by difference equations with positive coefficients*, J. Math. Phys. 31, 253–259, 1953

[Ma] V. G. Maz'ya, *Sobolev spaces*, Springer, N.Y., 1985

[RM] R. D. Richtmyer and K. W. Morton, *Difference Methods for Initial-Value Problems, 2nd ed.*, Interscience, New York, 1967

[LR3] M. Rogina and N. Limić, *Monotone numerical schemes for a Dirichlet problem for elliptic operators in divergence form*, in ICNAAM–2006 extended abstracts, 279–282, Wiley-VCH, 2006

[Sh] I. A. Shishmarev, *An introduction to the theory of elliptic equations (in Russian)*, Moscow Univ., 1979

[SMMM] A. A. Samarskii, P. P. Matus, V. I. Mazhukin and I. E. Mozolevski, *Monotone Difference Schemes for Equations with Mixed Derivatives*, Comput. Math. Appl. **44**, 501–510, 2002

[Ste] F. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, New Jersey, 1970

[Str] J. C. Strikwerda, *Finite Difference Schemes and Partial Differential Equations*, Wadsworth & Brooks/Cole, 1989

[Yo] D. M. Young, *Iterative Methods for Solving Partial Difference Equations of Elliptic Type*, thesis, Harvard University, Cambridge, Mass, 1950